

Highlighted sections are also highlighted in main printed

605

No

Zhang 10/728.665

10/728.665

28/09/2006

SEARCH HISTORY

=> d his ful

(FILE 'HOME' ENTERED AT 15:11:25 ON 28 SEP 2006)

FILE 'HCAPLUS' ENTERED AT 15:11:33 ON 28 SEP 2006

E MEDICHERLA SATYANARAYANA/AU

L1 29 SEA ABB=ON ("MEDICHERLA SATYA"/AU OR "MEDICHERLA SATYANANYANA"/AU OR "MEDICHERLA SATYANARAYANA"/AU)

E PROTTER ANDREW A/AU

L2 73 SEA ABB=ON ("PROTTER A A"/AU OR "PROTTER ANDREW"/AU OR "PROTTER ANDREW A"/AU OR "PROTTER ANDREW ASHER"/AU OR "PROTTER ANDY"/AU)

E SCHREINER GEORGE F/AU

L3 170 SEA ABB=ON ("SCHREINER GEORGE"/AU OR "SCHREINER GEORGE E"/AU OR "SCHREINER GEORGE F"/AU OR "SCHREINER GEORGE FREDERIC"/AU)

L4 5 SEA ABB=ON L1 AND L2 AND L3

L5 3 SEA ABB=ON L4 AND ?DIABETES?

SELECT RN L5 1-3

FILE 'REGISTRY' ENTERED AT 15:13:00 ON 28 SEP 2006

L6 22 SEA ABB=ON (165245-96-5/BI OR 50-99-7/BI OR 9004-10-8/BI OR 152060-53-2/BI OR 1670-87-7/BI OR 176023-64-6/BI OR 179800-23-8/BI OR 192333-55-4/BI OR 253-82-7/BI OR 309913-28-8/BI OR 309913-29-9/BI OR 309913-41-5/BI OR 309913-51-7/BI OR 309913-59-5/BI OR 309913-92-6/BI OR 309914-09-8/BI OR 309914-17-8/BI OR 309914-25-8/BI OR 309914-66-7/BI OR 309914-79-2/BI OR 309914-94-1/BI OR 627536-09-8/BI)

FILE 'HCAPLUS' ENTERED AT 15:13:18 ON 28 SEP 2006

L7 3 SEA ABB=ON L5 AND L6 *Inventor Search*

Compd Search FILE 'REGISTRY' ENTERED AT 15:15:48 ON 28 SEP 2006

L8 1 SEA ABB=ON 309913-51-7/RN

L9 1 SEA ABB=ON 309913-51-7/RN *Requested compd*

FILE 'HCAPLUS' ENTERED AT 15:16:44 ON 28 SEP 2006

L10 8 SEA ABB=ON L9

L11 1 SEA ABB=ON L10 AND ?DIABETES?

L12 8 SEA ABB=ON L10 OR L11

L13 6 SEA ABB=ON L12 AND (PRD<20031206 OR PD<20031206)

FILE 'MEDLINE, BIOSIS, EMBASE, JAPIO, JICST-EPLUS' ENTERED AT 15:17:49 ON 28 SEP 2006

L14 0 SEA ABB=ON L12

FILE 'USPATFULL' ENTERED AT 15:18:13 ON 28 SEP 2006

L15 5 SEA ABB=ON L12 AND (PRD<20031206 OR PD<20031206)

FILE 'HCAPLUS, USPATFULL' ENTERED AT 15:18:45 ON 28 SEP 2006

L16 9 DUP REMOV L13 L15 (2 DUPLICATES REMOVED) *9 hits for CAPLUS & usPatfull*

Text Search FILE 'REGISTRY' ENTERED AT 15:39:12 ON 28 SEP 2006

E P38 MAPK INHIBITOR/CN

L17 2 SEA ABB=ON ("P38 MAPK"/CN OR "P38 MAPK (BIOMPHALARIA GLABRATA STRAIN BS90 HEMOCYTE)"/CN OR "P38 MAPK INHIBITOR"/CN)

FILE 'HCAPLUS' ENTERED AT 15:39:42 ON 28 SEP 2006

L18 10002 SEA ABB=ON L17 OR P38(W)?MAPK?(W)?INHIBIT?

L19 320 SEA ABB=ON L18 AND ?DIABETES?
L20 25 SEA ABB=ON L19 AND (TYPE(W) (1 OR I) (W)?DIABETES?)
L21 6 SEA ABB=ON L20 AND (PRD<20021206 OR PD<20021206)

FILE 'MEDLINE, BIOSIS, EMBASE, JAPIO, JICST-EPLUS' ENTERED AT 15:41:28 ON 28 SEP 2006

L22 23 SEA ABB=ON L20 *23 cits from*

FILE 'USPATFULL' ENTERED AT 15:42:04 ON 28 SEP 2006

L23 28 SEA ABB=ON L20 AND (PRD<20021206 OR PD<20021206)

FILE 'USPATFULL, HCAPLUS' ENTERED AT 15:43:04 ON 28 SEP 2006

L24 34 DUP REMOV L23 L21 (0 DUPLICATES REMOVED) *34 cits from*

FILE HOME

FILE HCAPLUS

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 28 Sep 2006 VOL 145 ISS 14
FILE LAST UPDATED: 27 Sep 2006 (20060927/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

FILE REGISTRY

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 27 SEP 2006 HIGHEST RN 909000-49-3
DICTIONARY FILE UPDATES: 27 SEP 2006 HIGHEST RN 909000-49-3

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH June 30, 2006

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/ONLINE/UG/regprops.html>

FILE MEDLINE

FILE LAST UPDATED: 27 Sep 2006 (20060927/UP). FILE COVERS 1950 TO DATE.

On December 11, 2005, the 2006 MeSH terms were loaded.

The MEDLINE reload for 2006 is now (26 Feb.) available. For details on the 2006 reload, enter HELP RLOAD at an arrow prompt (=>).
See also:

<http://www.nlm.nih.gov/mesh/>
http://www.nlm.nih.gov/pubs/techbull/nd04/nd04_mesh.html
http://www.nlm.nih.gov/pubs/techbull/nd05/nd05_med_data_changes.html
http://www.nlm.nih.gov/pubs/techbull/nd05/nd05_2006_MeSH.html

OLDMEDLINE is covered back to 1950.

MEDLINE thesauri in the /CN, /CT, and /MN fields incorporate the MeSH 2006 vocabulary.

This file contains CAS Registry Numbers for easy and accurate substance identification.

FILE BIOSIS

FILE COVERS 1969 TO DATE.

CAS REGISTRY NUMBERS AND CHEMICAL NAMES (CNs) PRESENT
FROM JANUARY 1969 TO DATE.

RECORDS LAST ADDED: 27 September 2006 (20060927/ED)

FILE EMBASE

FILE COVERS 1974 TO 28 Sep 2006 (20060928/ED)

EMBASE has been reloaded. Enter HELP RLOAD for details.

EMBASE is now updated daily. SDI frequency remains weekly (default) and biweekly.

This file contains CAS Registry Numbers for easy and accurate substance identification.

FILE JAPIO

FILE LAST UPDATED: 3 APR 2006 <20060403/UP>

FILE COVERS APRIL 1973 TO DECEMBER 22, 2005

>>> GRAPHIC IMAGES AVAILABLE <<<

>>> NEW IPC8 DATA AND FUNCTIONALITY NOT YET AVAILABLE IN THIS FILE.
USE IPC7 FORMAT FOR SEARCHING THE IPC. WATCH THIS SPACE FOR FURTHER
DEVELOPMENTS AND SEE OUR NEWS SECTION FOR FURTHER INFORMATION
ABOUT THE IPC REFORM <<<

FILE JICST-EPLUS

FILE COVERS 1985 TO 26 SEP 2006 (20060926/ED)

THE JICST-EPLUS FILE HAS BEEN RELOADED TO REFLECT THE 1999 CONTROLLED
TERM (/CT) THESAURUS RELOAD.

FILE USPATFULL

FILE COVERS 1971 TO PATENT PUBLICATION DATE: 28 Sep 2006 (20060928/PD)

FILE LAST UPDATED: 28 Sep 2006 (20060928/ED)

HIGHEST GRANTED PATENT NUMBER: US7114185

HIGHEST APPLICATION PUBLICATION NUMBER: US2006218687

CA INDEXING IS CURRENT THROUGH 28 Sep 2006 (20060928/UPCA)
ISSUE CLASS FIELDS (/INCL) CURRENT THROUGH: 28 Sep 2006 (20060928/PD)
REVISED CLASS FIELDS (/NCL) LAST RELOADED: Jun 2006
USPTO MANUAL OF CLASSIFICATIONS THESAURUS ISSUE DATE: Jun 2006

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:sssptal700mjr

PASSWORD:

* * * * * RECONNECTED TO STN INTERNATIONAL * * * * *
SESSION RESUMED IN FILE 'MEDLINE, BIOSIS, EMBASE, JAPIO, JICST-EPLUS'
AT 15:52:54 ON 28 SEP 2006
FILE 'MEDLINE' ENTERED AT 15:52:54 ON 28 SEP 2006
FILE 'BIOSIS' ENTERED AT 15:52:54 ON 28 SEP 2006
Copyright (c) 2006 The Thomson Corporation
FILE 'EMBASE' ENTERED AT 15:52:54 ON 28 SEP 2006
Copyright (c) 2006 Elsevier B.V. All rights reserved.
FILE 'JAPIO' ENTERED AT 15:52:54 ON 28 SEP 2006
COPYRIGHT (C) 2006 Japanese Patent Office (JPO)- JAPIO
FILE 'JICST-EPLUS' ENTERED AT 15:52:54 ON 28 SEP 2006
COPYRIGHT (C) 2006 Japan Science and Technology Agency (JST)

INVENTOR SEARCH

=> d ibib abs hitstr l7 1-3

L7 ANSWER 1 OF 3 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2006:658530 HCAPLUS

DOCUMENT NUMBER: 145:262961

TITLE: Preventive and therapeutic potential of p38 α -selective mitogen-activated protein kinase inhibitor in nonobese diabetic mice with type 1 diabetes

AUTHOR(S): Medicherla, Satyanarayana; Protter, Andrew A.; Ma, Jing Ying; Mangadu, Ruban; Almirez, Ramona; Koppelman, Bruce; Kerr, Irene; Navas, Tony A.; Movius, Fabiola; Reddy, Mamatha; Liu, Yu-Wang; Luedtke, Gregory; Perumattam, John; Mavunkel, Babu; Dugar, Sundee; Schreiner, George F.

CORPORATE SOURCE: Scios Inc., Fremont, CA, USA

SOURCE: Journal of Pharmacology and Experimental Therapeutics (2006), 318(1), 99-107

CODEN: JPETAB; ISSN: 0022-3565

PUBLISHER: American Society for Pharmacology and Experimental Therapeutics

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Mitogen-activated protein kinases (MAPKs) and heat shock proteins (HSPs) are ubiquitous proteins that function within T cells in both normal and stress-related pathophysiol. states, including type 1 diabetes. The nonobese diabetic (NOD) mouse spontaneously develops T cell-mediated autoimmune pancreatic beta cell destruction that is similar to type 1 diabetes in humans. Because p38 MAPKs have been shown to modulate T cell function, we studied the effects of a p38 α MAPK-selective inhibitor, indole-5-carboxamide (SD-169), on the development and progression of type 1 diabetes in the NOD mouse. In preventive treatment studies, SD-169 significantly reduced p38 and HSP60 expression in T cells of the pancreatic beta islets. Following treatment, the incidence of diabetes as determined by blood glucose levels was significantly lower, and immunohistochem. of pancreatic beta islet tissue demonstrated significant reduction in CD5+ T cell infiltration in the SD-169 treatment group as compared with untreated NOD mice. In therapeutic studies using mildly and moderately hyperglycemic NOD mice, SD-169 treatment lowered blood glucose and improved glucose homeostasis. Furthermore, following cessation of SD-169 treatment, NOD mice showed significant arrest of diabetes. In conclusion, we report that this p38 α -selective inhibitor prevents the development and progression of diabetes in NOD mice by inhibiting T cell infiltration and activation, thereby preserving beta cell mass via inhibition of the p38 MAPK signaling pathway. These results have bearing on current prophylactic and therapeutic protocols using p38 α -selective inhibitors in the prediabetic period for children at high risk of type 1 diabetes, in the honeymoon period, and for adults with latent autoimmune diabetes.


IT 1670-87-7, SD 169

RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(SD 169; preventive and therapeutic potential of p38 α -selective MAPK inhibitor SD-169 in nonobese diabetic mice with type 1 diabetes)

RN 1670-87-7 HCAPLUS

CN 1H-Indole-5-carboxamide (9CI) (CA INDEX NAME)



IT 165245-96-5, p38 MAP kinase
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (preventive and therapeutic potential of p38 α -selective MAPK
 inhibitor SD-169 in nonobese diabetic mice with type 1 diabetes
)
 RN 165245-96-5 HCAPLUS
 CN Kinase (phosphorylating), protein, RK (9CI) (CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

REFERENCE COUNT: 37 THERE ARE 37 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 2 OF 3 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2004:546407 HCAPLUS

DOCUMENT NUMBER: 141:82337

TITLE: Treatment of obesity and associated conditions with
 TGF- β inhibitors

INVENTOR(S): Medicherla, Satyanarayana; Protter,
 Andrew A.; Schreiner, George F.

PATENT ASSIGNEE(S): Scios, Inc., USA

SOURCE: PCT Int. Appl., 89 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English


FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------|----------|-----------------|------------|
| WO 2004056352 | A1 | 20040708 | WO 2003-US40907 | 20031218 |
| W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, | | | | |
| CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, | | | | |
| GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, | | | | |
| LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, | | | | |
| PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, | | | | |
| UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW | | | | |
| RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, | | | | |
| BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, | | | | |
| ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, | | | | |
| TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG | | | | |
| CA 2513086 | AA | 20040708 | CA 2003-2513086 | 20031218 |
| AU 2003297460 | A1 | 20040714 | AU 2003-297460 | 20031218 |
| US 2004192583 | A1 | 20040930 | US 2003-742689 | 20031218 |
| EP 1589960 | A1 | 20051102 | EP 2003-813828 | 20031218 |
| R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, | | | | |
| IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK | | | | |
| JP 2006512369 | T2 | 20060413 | JP 2004-562359 | 20031218 |
| PRIORITY APPLN. INFO.: | | | US 2002-435856P | P 20021219 |
| | | | WO 2003-US40907 | W 20031218 |

OTHER SOURCE(S):
GI

MARPAT 141:82337




AB The invention concerns the treatment obesity and associated conditions with TGF- β inhibitors. More specifically, the invention concerns the use of TGF- β inhibitors in the treatment of obesity, type 2 diabetes, and pathol. conditions associated with obesity or type 2 diabetes. Quinazoline derivative I significantly restricted food intake and reduced the body weight of db/db obese mice. I also effectively modulated blood glucose levels.

IT 253-82-7D, Quinazoline, derivs.

RL: BSU (Biological study, unclassified); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(as TGF β inhibitor; TGF- β inhibitors for treatment of obesity and associated conditions)

RN 253-82-7 HCAPLUS

CN Quinazoline (6CI, 8CI, 9CI) (CA INDEX NAME)




IT 627536-09-8

RL: BSU (Biological study, unclassified); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(as TGF β -R1 inhibitor; TGF- β inhibitors for treatment of obesity and associated conditions)


RN 627536-09-8 HCAPLUS

CN 4-Pteridinamine, 2-(5-chloro-2-fluorophenyl)-N-4-pyridinyl- (9CI) (CA INDEX NAME)



IT 50-99-7, D-Glucose, biological studies
 RL: ADV (Adverse effect, including toxicity); BSU (Biological study, unclassified); BIOL (Biological study)
 (blood, treatment of damage caused to blood vessels, nerves and other internal structures by elevated levels of; TGF- β inhibitors for treatment of obesity and associated conditions)
 RN 50-99-7 HCAPLUS
 CN D-Glucose (8CI, 9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 152060-53-2, Type I TGF- β receptor kinase
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (inhibition of signaling through; TGF- β inhibitors for treatment of obesity and associated conditions)
 RN 152060-53-2 HCAPLUS
 CN Kinase (phosphorylating), β -transforming growth factor type I receptor (9CI) (CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

IT 9004-10-8, Insulin, biological studies
 RL: ADV (Adverse effect, including toxicity); BSU (Biological study, unclassified); BIOL (Biological study)
 (resistance, treatment of; TGF- β inhibitors for treatment of obesity and associated conditions)
 RN 9004-10-8 HCAPLUS
 CN Insulin (9CI) (CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***


REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 3 OF 3 HCAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 2004:515679 HCAPLUS
 DOCUMENT NUMBER: 141:47344

TITLE: Methods using p38 mitogen-activated protein kinase inhibitors for treating **diabetes**
 INVENTOR(S): **Medicherla, Satyanarayana; Protter, Andrew A.; Schreiner, George F.**
 PATENT ASSIGNEE(S): Scios Inc., USA
 SOURCE: PCT Int. Appl., 104 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------|----------|-----------------|------------|
| WO 2004053107 | A2 | 20040624 | WO 2003-US40140 | 20031205 |
| WO 2004053107 | A3 | 20041007 | | |
| W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW | | | | |
| RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG | | | | |
| CA 2511763 | AA | 20040624 | CA 2003-2511763 | 20031205 |
| AU 2003299652 | A1 | 20040630 | AU 2003-299652 | 20031205 |
| US 2004171659 | A1 | 20040902 | US 2003-728665 | 20031205 |
| EP 1583535 | A2 | 20051012 | EP 2003-799936 | 20031205 |
| R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK | | | | |
| JP 2006510654 | T2 | 20060330 | JP 2004-558236 | 20031205 |
| PRIORITY APPLN. INFO.: | | | | |
| | | | US 2002-431241P | P 20021206 |
| | | | WO 2003-US40140 | W 20031205 |
| AB The invention discloses methods for treating diabetes by administering p38 mitogen-activated protein kinase inhibitors. The invention also discloses methods of decreasing blood glucose level in diabetes patients by administering p38 mitogen-activated protein kinase inhibitors. | | | | |
| IT 50-99-7, D-Glucose, biological studies 9004-10-8, Insulin, biological studies 165245-96-5, p38 MAP kinase 176023-64-6, p38 γ MAP kinase 179800-23-8, p38 β Mitogen-activated protein kinase 192333-55-4, p38 δ MAP kinase | | | | |
| RL: BSU (Biological study, unclassified); BIOL (Biological study) (p38 MAP kinase inhibitors for treatment of diabetes) | | | | |
| RN 50-99-7 HCAPLUS | | | | |
| CN D-Glucose (8CI, 9CI) (CA INDEX NAME) | | | | |

Absolute stereochemistry.



RN 9004-10-8 HCAPLUS
CN Insulin (9CI) (CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

RN 165245-96-5 HCAPLUS
CN Kinase (phosphorylating), protein, RK (9CI) (CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

RN 176023-64-6 HCAPLUS
CN Kinase (phosphorylating), stress-activated protein, 3 (9CI) (CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

RN 179800-23-8 HCAPLUS
CN Kinase (phosphorylating), protein p38 β (9CI) (CA INDEX NAME)


*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

RN 192333-55-4 HCAPLUS
CN Kinase (phosphorylating), protein, SAPK4 (9CI) (CA INDEX NAME)


*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

IT 309913-28-8 309913-29-9 309913-41-5
309913-51-7 309913-59-5 309913-92-6
309914-09-8 309914-17-8 309914-25-8
309914-66-7 309914-79-2 309914-94-1
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL
(Biological study); USES (Uses)
(p38 MAP kinase inhibitors for treatment of **diabetes**)


RN 309913-28-8 HCAPLUS
CN 1H-Indole-3-acetic acid, 5-[[4-[(4-fluorophenyl)methyl]-1-piperidinyl]carbonyl]-6-methoxy-1-methyl- α -oxo- (9CI) (CA INDEX NAME)




RN 309913-29-9 HCAPLUS
CN Piperazine, 1-[[5-[[4-[(4-fluorophenyl)methyl]-1-piperidinyl]carbonyl]-6-methoxy-1-methyl-1H-indol-3-yl]oxoacetyl]-4-methyl- (9CI) (CA INDEX NAME)




RN 309913-41-5 HCAPLUS
CN 1H-Indole-3-acetamide, 5-[[4-[(4-fluorophenyl)methyl]-2,5-dimethyl-1-piperazinyl]carbonyl]-6-methoxy-N,N,1-trimethyl- α -oxo- (9CI) (CA INDEX NAME)




RN 309913-51-7 HCAPLUS
CN 1H-Indole-3-acetamide, 5-[[4-[(4-fluorophenyl)methyl]-1-piperidinyl]carbonyl]-6-methoxy-N,N,1-trimethyl- α -oxo- (9CI) (CA INDEX NAME)



RN 309913-59-5 HCAPLUS
CN 1H-Indole-3-acetamide, 6-chloro-5-[[4-[(4-fluorophenyl)methyl]-2,5-dimethyl-1-piperazinyl]carbonyl]-N,N,1-trimethyl- α -oxo- (9CI) (CA INDEX NAME)




RN 309913-92-6 HCAPLUS
CN 1H-Indole-3-acetamide, 5-[[4-[(4-fluorophenyl)methylene]-1-piperidinyl]carbonyl]-6-methoxy-N,N,1-trimethyl- α -oxo- (9CI) (CA INDEX NAME)



RN 309914-09-8 HCAPLUS


CN Piperidine, 1-[[6-chloro-1-methyl-3-(oxo-1-pyrrolidinylacetyl)-1H-indol-5-yl]carbonyl]-4-[(4-fluorophenyl)methyl]- (9CI) (CA INDEX NAME)



RN 309914-17-8 HCAPLUS

CN 1H-Indole-3-acetamide, 5-[[[(2R,5S)-4-[(4-fluorophenyl)methyl]-2,5-dimethyl-1-piperazinyl]carbonyl]-6-methoxy-1-(methoxymethyl)-N,N-dimethyl-α-oxo- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 309914-25-8 HCAPLUS


CN Morpholine, 4-[[5-[[[(2R,5S)-4-[(4-fluorophenyl)methyl]-2,5-dimethyl-1-piperazinyl]carbonyl]-6-methoxy-1-methyl-1H-indol-3-yl]oxoacetyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 309914-66-7 HCAPLUS

CN Morpholine, 4-[[5-[[4-[(4-fluorophenyl)methyl]-1-piperidinyl]carbonyl]-6-methoxy-1H-indol-3-yl]oxoacetyl]- (9CI) (CA INDEX NAME)




RN 309914-79-2 HCAPLUS

CN 1H-Indole-3-acetamide, 5-[[4-[(4-fluorophenyl)methyl]-2,5-dimethyl-1-piperazinyl]carbonyl]-6-methoxy-N,N-dimethyl-α-oxo- (9CI) (CA INDEX NAME)



RN 309914-94-1 HCAPLUS


CN Piperazine, 1-[[6-chloro-5-[[4-[(4-fluorophenyl)methyl]-1-piperidinyl]carbonyl]-2-methyl-1H-indol-3-yl]oxoacetyl]-4-methyl- (9CI) (CA INDEX NAME)



DISPLAY OF ELECTED SPECIE

=> d

L8 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2006 ACS on STN
RN 309913-51-7 REGISTRY
ED Entered STN: 20 Dec 2000
CN 1H-Indole-3-acetamide, 5-[[4-[(4-fluorophenyl)methyl]-1-piperidinyl]carbonyl]-6-methoxy-N,N,1-trimethyl- α -oxo- (9CI) (CA INDEX NAME)
MF C27 H30 F N3 O4
SR CA
LC STN Files: CA, CAPLUS, PROUSDDR, SYNTHLINE, TOXCENTER, USPATFULL



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

8 REFERENCES IN FILE CA (1907 TO DATE)

8 REFERENCES IN FILE CAPLUS (1907 TO DATE)

ED Entered STN: 20 Dec 2000

COMPOUND SEARCH - CAPLUS & USPATFULL SEARCH

=> d que stat l16

L9 1 SEA FILE=REGISTRY ABB=ON 309913-51-7/RN
L10 8 SEA FILE=HCAPLUS ABB=ON L9
L11 1 SEA FILE=HCAPLUS ABB=ON L10 AND ?DIABETES?
L12 8 SEA FILE=HCAPLUS ABB=ON L10 OR L11
L13 6 SEA FILE=HCAPLUS ABB=ON L12 AND (PRD<20031206 OR PD<20031206)
L15 5 SEA FILE=USPATFULL ABB=ON L12 AND (PRD<20031206 OR PD<20031206)
L16 9 DUP REMOV L13 L15 (2 DUPLICATES REMOVED)

=> d ibib abs hitstr l16 1-9

L16 ANSWER 1 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN DUPLICATE 1

ACCESSION NUMBER: 2005:162022 HCAPLUS

DOCUMENT NUMBER: 142:254591

TITLE: Methods of screening for compounds that selectively inhibit p38 MAP kinase α isoenzymes for use as immunomodulators

INVENTOR(S): Kirschenbaum, Ford; Higgins, Linda S.; Schreiner, George F.

PATENT ASSIGNEE(S): USA

SOURCE: U.S. Pat. Appl. Publ., 41 pp., Cont.-in-part of U.S. Ser. No. 683,656.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------------------|------|----------|-----------------|-----------------|
| US 2005043212 | A1 | 20050224 | US 2004-830834 | 20040422 <-- |
| US 2004176598 | A1 | 20040909 | US 2003-683656 | 20031009 <-- |
| PRIORITY APPLN. INFO.: | | | US 2002-417599P | P 20021009 <-- |
| | | | US 2003-683656 | A2 20031009 <-- |


AB The invention relates to methods of screening for compds. that selectively inhibit p38 MAP kinase α isoenzymes for use as immunomodulators. Inhibitors of p38 MAP kinase α isoenzyme include siRNA and SB203580.

IT 309913-51-7

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(methods of screening for compds. that selectively inhibit p38 MAP kinase α isoenzymes for use as immunomodulators)



RN 309913-51-7 HCAPLUS

CN 1H-Indole-3-acetamide, 5-[[4-[(4-fluorophenyl)methyl]-1-piperidinyl]carbonyl]-6-methoxy-N,N,1-trimethyl- α -oxo- (9CI) (CA INDEX NAME)



L16 ANSWER 2 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN DUPLICATE 2
 ACCESSION NUMBER: 2005:232421 HCAPLUS
 DOCUMENT NUMBER: 142:316692
 TITLE: Preparation of indolylcarboxamide derivatives as
 inhibitors of p38 kinase
 INVENTOR(S): Mavunkel, Babu J.; Chakravarty, Sarvajit; Perumattam,
 John J.; Dugar, Sundeeep; Lu, Qing; Liang, Xi
 PATENT ASSIGNEE(S): Scios, Inc., USA
 SOURCE: U.S., 65 pp., Cont.-in-part of U.S. 6,589,954.
 CODEN: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 7
 PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------------------|-------------------|----------|-----------------|-----------------|
| US 6867209 | B1 | 20050315 | US 2000-575060 | 20000519 <-- |
| US 6130235 | A | 20001010 | US 1998-128137 | 19980803 <-- |
| US 6340685 | B1 | 20020122 | US 1999-275176 | 19990324 <-- |
| US 6589954 | B1 | 20030708 | US 1999-316761 | 19990521 <-- |
| US 2003158417 | A1 | 20030821 | US 2002-146703 | 20020514 <-- |
| US 2003144520 | A1 | 20030731 | US 2002-157048 | 20020528 <-- |
| US 6864260 | B2 | 20050308 | | |
| US 2003162970 | A1 | 20030828 | US 2002-156996 | 20020528 <-- |
| US 2003195355 | A1 | 20031016 | US 2002-156997 | 20020528 <-- |
| PRIORITY APPLN. INFO.: | | | US 1998-86531P | P 19980522 <-- |
| | | | US 1998-128137 | A2 19980803 <-- |
| | | | US 1999-275176 | A2 19990324 <-- |
| | | | US 1999-316761 | A2 19990521 <-- |
| | | | US 1999-154594P | P 19990917 <-- |
| | | | US 2000-202608P | P 20000509 <-- |
| | | | US 2000-575060 | A1 20000519 <-- |
| OTHER SOURCE(S): | MARPAT 142:316692 | | | |
| GI | | | | |



AB Title compds. I [X independently = CA, CR4A, CR5, CR52, NR6, or N; L1 = CO, SO2, or alkylene; L2 = (un)substituted-alkylene or -alkenylene; Ar = (un)substituted aryl group with substituents consisting of alkyl, alkenyl, halo, CN, etc.; Z = N or CR7 wherein R7 = H or non-interfering substituent; R1 = H, alkyl, alkenyl, alkynyl, aryl, arylalkyl, etc.; R2 independently = halo, alkyl, OH, alkoxy, etc.; R3 independently = CN, CF3, NO2, alkyl, aryl, acyl, etc.; R4 = H, halo, alkyl or alkenyl; R5 independently = H, halo, alkyl, OH, etc.; R6 = H, alkyl, alkenyl, aryl, acyl, aroyl, etc.; A = -WiCOXjY wherein Y is COR8 wherein R8 = H, (un)substituted-alkyl, -alkenyl, -alkynyl, etc.; W and X = (un)substituted-alkylene, -alkenylene, -alkynylene; Y = tetrazole, 1,2,3-triazole, 1,2,4-triazole, or imidazole and each of i and j independently = 0 or 1; m = 0-4; n = 0-3], and their pharmaceutically acceptable salts are prepared and disclosed as useful for treatment of rheumatoid arthritis. Thus, e.g., II, was prepared by carbonylation of 6-methoxy-(4-benzylpiperidinyl)-indole-5-carboxamide with oxalyl chloride and subsequent amination using 4-methylpiperazine. ELISA assays for evaluation of inhibition of p38 kinase by I revealed that all compds. of the invention possessed IC50 values in the range of 0.1-1.5 μ M. I as inhibitors of p38 kinase should prove useful in the treatment of rheumatoid arthritis.


IT 309913-51-7P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of indolylcarboxamide derivs. as p38 kinase inhibitors)

RN 309913-51-7 HCAPLUS

CN 1H-Indole-3-acetamide, 5-[[4-[(4-fluorophenyl)methyl]-1-piperidinyl]carbonyl]-6-methoxy-N,N,1-trimethyl- α -oxo- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 54 THERE ARE 54 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L16 ANSWER 3 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2004:515679 HCAPLUS

DOCUMENT NUMBER: 141:47344

TITLE: Methods using p38 mitogen-activated protein kinase inhibitors for treating **diabetes**

INVENTOR(S): Medicherla, Satyanarayana; Protter, Andrew A.; Schreiner, George F.

PATENT ASSIGNEE(S): Scios Inc., USA

SOURCE: PCT Int. Appl., 104 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------------------|--|----------|-----------------|----------------|
| WO 2004053107 | A2 | 20040624 | WO 2003-US40140 | 20031205 <-- |
| WO 2004053107 | A3 | 20041007 | | |
| W: | AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW | | | |
| RW: | BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG | | | |
| CA 2511763 | AA | 20040624 | CA 2003-2511763 | 20031205 <-- |
| AU 2003299652 | A1 | 20040630 | AU 2003-299652 | 20031205 <-- |
| US 2004171659 | A1 | 20040902 | US 2003-728665 | 20031205 <-- |
| EP 1583535 | A2 | 20051012 | EP 2003-799936 | 20031205 <-- |
| R: | AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK | | | |
| JP 2006510654 | T2 | 20060330 | JP 2004-558236 | 20031205 <-- |
| PRIORITY APPLN. INFO.: | | | US 2002-431241P | P 20021206 <-- |
| | | | WO 2003-US40140 | W 20031205 <-- |


AB The invention discloses methods for treating **diabetes** by administering p38 mitogen-activated protein kinase inhibitors. The invention also discloses methods of decreasing blood glucose level in **diabetes** patients by administering p38 mitogen-activated protein kinase inhibitors.

IT 309913-51-7

RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(p38 MAP kinase inhibitors for treatment of diabetes)

RN 309913-51-7 HCAPLUS

CN 1H-Indole-3-acetamide, 5-[[4-[(4-fluorophenyl)methyl]-1-piperidinyl]carbonyl]-6-methoxy-N,N,1-trimethyl- α -oxo- (9CI) (CA INDEX NAME)

L16 ANSWER 4 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2004:203623 HCAPLUS

DOCUMENT NUMBER: 140:247108

TITLE: Bone healing and promoting osteogenesis by administration of a p38 MAP kinase inhibitor

INVENTOR(S): Protter, Andrew Asher; Liu, David Y.

PATENT ASSIGNEE(S): Scios Inc., USA

SOURCE: PCT Int. Appl., 127 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------|----------|-----------------|----------------|
| WO 2004019873 | A2 | 20040311 | WO 2003-US26839 | 20030829 <-- |
| WO 2004019873 | C2 | 20040624 | | |
| WO 2004019873 | A3 | 20041007 | | |
| W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW | | | | |
| RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG | | | | |
| CA 2497240 | AA | 20040311 | CA 2003-2497240 | 20030829 <-- |
| AU 2003262911 | A1 | 20040319 | AU 2003-262911 | 20030829 <-- |
| US 2004162289 | A1 | 20040819 | US 2003-651934 | 20030829 <-- |
| EP 1539121 | A2 | 20050615 | EP 2003-791848 | 20030829 <-- |
| R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK | | | | |
| JP 2006508051 | T2 | 20060309 | JP 2004-531559 | 20030829 <-- |
| PRIORITY APPLN. INFO.: | | | US 2002-406664P | P 20020829 <-- |
| | | | WO 2003-US26839 | W 20030829 <-- |

OTHER SOURCE(S): MARPAT 140:247108


AB The invention discloses methods of bone healing by administering a p38 MAP kinase inhibitor. Specifically, the invention provides methods of

treating bone fractures, bone diseases, bone grafting, especially enhancing bone healing following facial reconstruction, maxillary reconstruction, mandibular reconstruction or tooth extraction, enhancing long bone extension, enhancing prosthetic ingrowth, and increasing bone synostosis by administering a p38 MAP kinase inhibitor.

IT 309913-51-7
 RL: ADV (Adverse effect, including toxicity); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (bone healing and promoting osteogenesis by administration of a p38 MAP kinase inhibitor)

RN 309913-51-7 HCAPLUS

CN 1H-Indole-3-acetamide, 5-[[4-[(4-fluorophenyl)methyl]-1-piperidinyl]carbonyl]-6-methoxy-N,N,1-trimethyl- α -oxo- (9CI) (CA INDEX NAME)



L16 ANSWER 5 OF 9 USPATFULL on STN

ACCESSION NUMBER: 2004:221887 USPATFULL

TITLE: Methods for treating **diabetes**

INVENTOR(S): Medicherla, Satyanarayana, Cupertino, CA, UNITED STATES
 Protter, Andrew A., Palo Alto, CA, UNITED STATES
 Schreiner, George F., Los Altos, CA, UNITED STATES

| | NUMBER | KIND | DATE |
|---------------------|----------------|------|---------------|
| PATENT INFORMATION: | US 2004171659 | A1 | 20040902 |
| APPLICATION INFO.: | US 2003-728665 | A1 | 20031205 (10) |

| | NUMBER | DATE |
|-----------------------|---|-------------------|
| PRIORITY INFORMATION: | US 2002-431241P | 20021206 (60) <-- |
| DOCUMENT TYPE: | Utility | |
| FILE SEGMENT: | APPLICATION | |
| LEGAL REPRESENTATIVE: | MORRISON & FOERSTER LLP, 3811 VALLEY CENTRE DRIVE, SUITE 500, SAN DIEGO, CA, 92130-2332 | |
| NUMBER OF CLAIMS: | 11 | |
| EXEMPLARY CLAIM: | 1 | |
| NUMBER OF DRAWINGS: | 10 Drawing Page(s) | |
| LINE COUNT: | 3428 | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB The invention is directed to methods of treating **diabetes** by administering p38 mitogen activated protein kinase inhibitors. The invention is also directed to methods of decreasing blood glucose level in **diabetes** patients by administering p38 mitogen activated protein kinase inhibitors.


CAS INDEXING IS AVAILABLE FOR THIS PATENT.

IT 309913-51-7

(p38 MAP kinase inhibitors for treatment of diabetes)

RN 309913-51-7 USPTAFULL

CN 1H-Indole-3-acetamide, 5-[[4-[(4-fluorophenyl)methyl]-1-piperidinyl]carbonyl]-6-methoxy-N,N,1-trimethyl- α -oxo- (9CI) (CA INDEX NAME)



L16 ANSWER 6 OF 9 USPTAFULL on STN

ACCESSION NUMBER: 2004:209855 USPTAFULL

TITLE: Methods of promoting osteogenesis

INVENTOR(S): Protter, Andrew A., Palo Alto, CA, UNITED STATES
Liu, David Y., Palo Alto, CA, UNITED STATES

| | NUMBER | KIND | DATE |
|---------------------|----------------|------|---------------|
| PATENT INFORMATION: | US 2004162289 | A1 | 20040819 |
| APPLICATION INFO.: | US 2003-651934 | A1 | 20030829 (10) |

| | NUMBER | DATE |
|-----------------------|---|-------------------|
| PRIORITY INFORMATION: | US 2002-406664P | 20020829 (60) <-- |
| DOCUMENT TYPE: | Utility | |
| FILE SEGMENT: | APPLICATION | |
| LEGAL REPRESENTATIVE: | MORRISON & FOERSTER LLP, 3811 VALLEY CENTRE DRIVE, SUITE 500, SAN DIEGO, CA, 92130-2332 | |
| NUMBER OF CLAIMS: | 18 | |
| EXEMPLARY CLAIM: | 1 | |
| NUMBER OF DRAWINGS: | 13 Drawing Page(s) | |
| LINE COUNT: | 3708 | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB The invention is directed to methods of bone healing by administering a p38 MAP kinase inhibitor. The invention is directed to methods of treating bone fractures, bone diseases, bone grafting, especially enhancing bone healing following facial reconstruction, maxillary reconstruction, mandibular reconstruction or tooth extraction, enhancing long bone extension, enhancing prosthetic ingrowth, and increasing bone synostosis by administering a p38 MAP kinase inhibitor.


CAS INDEXING IS AVAILABLE FOR THIS PATENT.

IT 309913-51-7

(bone healing and promoting osteogenesis by administration of a p38 MAP kinase inhibitor)

RN 309913-51-7 USPTAFULL

CN 1H-Indole-3-acetamide, 5-[[4-[(4-fluorophenyl)methyl]-1-piperidinyl]carbonyl]-6-methoxy-N,N,1-trimethyl- α -oxo- (9CI) (CA INDEX NAME)



L16 ANSWER 7 OF 9 USPATFULL on STN

ACCESSION NUMBER: 2004:13467 USPATFULL

TITLE: Method to treat cystic fibrosis

INVENTOR(S): Higgins, Linda S., Palo Alto, CA, UNITED STATES

Liu, David Y., Palo Alto, CA, UNITED STATES

Protter, Andrew A., Palo Alto, CA, UNITED STATES

| | NUMBER | KIND | DATE |
|---------------------|----------------|------|---------------|
| PATENT INFORMATION: | US 2004009990 | A1 | 20040115 |
| APPLICATION INFO.: | US 2002-291243 | A1 | 20021108 (10) |

| | NUMBER | DATE |
|-----------------------|--|---------------|
| PRIORITY INFORMATION: | US 2001-338209P | 20011109 (60) |
| DOCUMENT TYPE: | Utility | |
| FILE SEGMENT: | APPLICATION | |
| LEGAL REPRESENTATIVE: | Kate H. Murashige, Morrison & Foerster LLP, Suite 500, 3811 Valley Centre Drive, San Diego, CA, 92130 | |
| NUMBER OF CLAIMS: | 40 | |
| EXEMPLARY CLAIM: | 1 | |
| LINE COUNT: | 1187 | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB The invention is directed to methods to treat cystic fibrosis by administering certain imidazole derivatives.


CAS INDEXING IS AVAILABLE FOR THIS PATENT.

IT 309913-51-7P

(imidazole derivs. for treatment of cystic fibrosis)

RN 309913-51-7 USPATFULL

CN 1H-Indole-3-acetamide, 5-[[4-[(4-fluorophenyl)methyl]-1-piperidinyl]carbonyl]-6-methoxy-N,N,1-trimethyl- α -oxo- (9CI) (CA INDEX NAME)



L16 ANSWER 8 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2003:396662 HCAPLUS

DOCUMENT NUMBER: 138:379271

TITLE: Method using imidazole derivatives to treat cystic fibrosis

INVENTOR(S): Higgins, Linda S.; Liu, David Y.; Protter, Andrew A.

PATENT ASSIGNEE(S): Scios Inc., USA

SOURCE: PCT Int. Appl., 42 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------|----------|-----------------|----------------|
| WO 2003041644 | A2 | 20030522 | WO 2002-US35939 | 20021108 <-- |
| WO 2003041644 | A3 | 20031113 | | |
| W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW | | | | |
| RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG | | | | |
| CA 2466665 | AA | 20030522 | CA 2002-2466665 | 20021108 <-- |
| US 2004009990 | A1 | 20040115 | US 2002-291243 | 20021108 <-- |
| EP 1453515 | A2 | 20040908 | EP 2002-778799 | 20021108 <-- |
| R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK | | | | |
| BR 2002014020 | A | 20041013 | BR 2002-14020 | 20021108 <-- |
| TR 200401028 | T2 | 20041122 | TR 2004-1028 | 20021108 <-- |
| JP 2005511616 | T2 | 20050428 | JP 2003-543531 | 20021108 <-- |
| PRIORITY APPLN. INFO.: | | | US 2001-338209P | P 20011109 <-- |
| | | | WO 2002-US35939 | W 20021108 <-- |

OTHER SOURCE(S): MARPAT 138:379271

AB The invention is directed to methods to treat cystic fibrosis by administering certain imidazole derivs.


IT 309913-51-7P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(imidazole derivs. for treatment of cystic fibrosis)

RN 309913-51-7 HCAPLUS

CN 1H-Indole-3-acetamide, 5-[[4-[(4-fluorophenyl)methyl]-1-piperidinyl]carbonyl]-6-methoxy-N,N,1-trimethyl- α -oxo- (9CI) (CA INDEX NAME)



L16 ANSWER 9 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2000:842127 HCAPLUS

DOCUMENT NUMBER: 134:17503

TITLE: Preparation of 5-[4-benzylpiperidinyl(piperazinyl)]-indolecarboxamides as inhibitors of p38 kinase

INVENTOR(S): Mavunkel, Babu J.; Chakravarty, Sarvajit; Perumattam, John J.; Dugar, Sundeeep; Lu, Qing; Liang, Xi

PATENT ASSIGNEE(S): Scios Inc., USA

SOURCE: PCT Int. Appl., 85 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 7



PATENT INFORMATION:

vs Pat 6,864,260

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------|----------|-----------------|-----------------|
| WO 2000071535 | A1 | 20001130 | WO 2000-US14003 | 20000519 <-- |
| W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW | | | | |
| RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG | | | | |
| US 6589954 | B1 | 20030708 | US 1999-316761 | 19990521 <-- |
| CA 2372567 | AA | 20001130 | CA 2000-2372567 | 20000519 <-- |
| EP 1178983 | A1 | 20020213 | EP 2000-939322 | 20000519 <-- |
| R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO | | | | |
| BR 2000011274 | A | 20020226 | BR 2000-11274 | 20000519 <-- |
| NZ 515285 | A | 20040130 | NZ 2000-515285 | 20000519 <-- |
| AU 772295 | B2 | 20040422 | AU 2000-54424 | 20000519 <-- |
| RU 2278115 | C2 | 20060620 | RU 2001-134501 | 20000519 <-- |
| BG 106091 | A | 20020628 | BG 2001-106091 | 20011108 <-- |
| HR 2001000854 | A1 | 20030430 | HR 2001-854 | 20011119 <-- |
| NO 2001005655 | A | 20020118 | NO 2001-5655 | 20011120 <-- |
| AU 2004203356 | A1 | 20040819 | AU 2004-203356 | 20040722 <-- |
| PRIORITY APPLN. INFO.: | | | US 1999-316761 | A 19990521 <-- |
| | | | US 1999-154594P | P 19990917 <-- |
| | | | US 2000-202608P | P 20000509 <-- |
| | | | US 1998-86531P | P 19980522 <-- |
| | | | US 1998-128137 | A2 19980803 <-- |
| | | | US 1999-275176 | A2 19990324 <-- |
| | | | WO 2000-US14003 | W 20000519 <-- |

OTHER SOURCE(S): MARPAT 134:17503

GI



AB The title compds. [I; one Z2 = CA, CR8A and the other = CR1, CR12, NR6, N (wherein R1, R6, R8 = H, noninterfering substituent; A = WiCOXjY; Y = COR2, an isostere; R2 = H, noninterfering substituent; W, X = spacer of 2-6Å; i, j = 0-1); Z3 = NR7, O; R3 = noninterfering substituent; n = 0-3; L1, L2 = linker; R4 = noninterfering substituent; m = 0-4; Z1 = CR5, N (R5 = H, noninterfering substituent); l, k = 0-2, wherein the sum of l and k = 0-3; Ar = aryl substituted with 0-5 noninterfering substituents, wherein two noninterfering substituents can form a fused ring; the distance between the atom of Ar linked to L2 and the center of the α ring is 4.5-24Å] which inhibit p38- α kinase (biol. data given), were prepared Thus, treating 6-methoxy-(4-benzylpiperidinyl)-indole-5-carboxamide with oxalyl chloride in CH2Cl2 afforded the indole-5-carboxamide II.


IT 309913-51-7P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

```
(preparation of 5-[4-benzylpiperidinyl(piperazinyl)]-indolecarboxamides as
inhibitors of p38 kinase)
```

RN 309913-51-7 HCAPLUS

CN 1H-Indole-3-acetamide, 5-[[4-[(4-fluorophenyl)methyl]-1-piperidinyl]carbonyl]-6-methoxy-N,N,1-trimethyl- α -oxo- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

TEXT SEARCH - CAPLUS & USPATFULL

=> d que stat l24

L17 2 SEA FILE=REGISTRY ABB=ON ("P38 MAPK"/CN OR "P38 MAPK (BIOMPHAL
ARIA GLABRATA STRAIN BS90 HEMOCYTE)"/CN OR "P38 MAPK INHIBITOR"
/CN)
L18 10002 SEA FILE=HCAPLUS ABB=ON L17 OR P38(W)?MAPK?(W)?INHIBIT?
L19 320 SEA FILE=HCAPLUS ABB=ON L18 AND ?DIABETES?
L20 25 SEA FILE=HCAPLUS ABB=ON L19 AND (TYPE(W)(1 OR I)(W)?DIABETES?)
L21 6 SEA FILE=HCAPLUS ABB=ON L20 AND (PRD<20021206 OR PD<20021206)
L23 28 SEA FILE=USPATFULL ABB=ON L20 AND (PRD<20021206 OR PD<20021206
)
L24 34 DUP REMOV L23 L21 (0 DUPLICATES REMOVED)

=> d ibib abs l24 1-34

L24 ANSWER 1 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2006:175374 USPATFULL
TITLE: Protein kinase inhibitors
INVENTOR(S): Burns, Christopher John, Melbourne, AUSTRALIA
Bu, Xianying, Rosanna East, AUSTRALIA
Wilks, Andrew Frederick, South Yarra, AUSTRALIA

| | NUMBER | KIND | DATE |
|-----------------------|--|------|---------------|
| PATENT INFORMATION: | US 2006148824 | A1 | 20060706 |
| APPLICATION INFO.: | US 2006-367248 | A1 | 20060302 (11) |
| RELATED APPLN. INFO.: | Continuation of Ser. No. US 2003-469303, filed on 29 Aug 2003, PENDING A 371 of International Ser. No. WO 2003-AU629, filed on 23 May 2003 | | |

| | NUMBER | DATE | |
|-----------------------|--|---------------|-----|
| PRIORITY INFORMATION: | AU 2002-200200000251520020523 | | <-- |
| | US 2002-399070P | 20020726 (60) | <-- |
| DOCUMENT TYPE: | Utility | | |
| FILE SEGMENT: | APPLICATION | | |
| LEGAL REPRESENTATIVE: | MORRISON & FOERSTER LLP, 12531 HIGH BLUFF DRIVE, SUITE 100, SAN DIEGO, CA, 92130-2040, US | | |
| NUMBER OF CLAIMS: | 24 | | |
| EXEMPLARY CLAIM: | 1 | | |
| NUMBER OF DRAWINGS: | 2 Drawing Page(s) | | |
| LINE COUNT: | 1671 | | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB A compound of the general formula: ##STR1## or pharmaceutically
acceptable salts, hydrates, solvates, crystal forms of diastereomers
thereof is described. Method of inhibiting a protein kinase using
compounds of Formula I are also described.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 2 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2006:22076 USPATFULL
TITLE: Methods for treating and preventing insulin resistance
and related disorders
INVENTOR(S): Greenberg, Andrew S., Boston, MA, UNITED STATES

| | NUMBER | KIND | DATE |
|-----------------------|---|------|---------------|
| PATENT INFORMATION: | US 2006019910 | A1 | 20060126 |
| APPLICATION INFO.: | US 2004-977328 | A1 | 20041029 (10) |
| RELATED APPLN. INFO.: | Continuation of Ser. No. US 2000-690647, filed on 17 Oct 2000, GRANTED, Pat. No. US 6897019 | | |
| | Continuation-in-part of Ser. No. WO 1999-US8364, filed on 16 Apr 1999, PENDING | | |

| | NUMBER | DATE | |
|-----------------------|---|---------------|-----|
| PRIORITY INFORMATION: | US 1998-82152P | 19980417 (60) | <-- |
| | US 1998-82741P | 19980423 (60) | <-- |
| DOCUMENT TYPE: | Utility | | |
| FILE SEGMENT: | APPLICATION | | |
| LEGAL REPRESENTATIVE: | FOLEY HOAG, LLP, PATENT GROUP, WORLD TRADE CENTER WEST, 155 SEAPORT BLVD, BOSTON, MA, 02110, US | | |
| NUMBER OF CLAIMS: | 37 | | |
| EXEMPLARY CLAIM: | 1-18 | | |
| NUMBER OF DRAWINGS: | 14 Drawing Page(s) | | |
| LINE COUNT: | 2759 | | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB The invention provides methods, therapeutics and kits for treating and preventing diseases or conditions associated with excessive lipolysis, in particular TNF- α induced lipolysis, and/or excessive free fatty acid levels. Exemplary conditions include insulin-resistance, **diabetes**, in particular NIDDM, obesity, glucose intolerance, hyperinsulinemia, polycystic ovary syndrome, and coronary artery disease. In a preferred embodiment, the method includes administering to a subject in need a pharmaceutically effective amount of an inhibitor of the JNK signal transduction pathway and/or an inhibitor of the MAPK/ERK signal transduction pathway.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 3 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2006:169991 USPATFULL

TITLE: Substituted pyrazoles as p38 kinase inhibitors

INVENTOR(S): Naraian, Ashok S., Ballwin, MO, UNITED STATES
 Clare, Michael, Skokie, IL, UNITED STATES
 Collins, Paul W., Deerfield, IL, UNITED STATES
 Crich, Joyce Zuowu, Glenview, IL, UNITED STATES
 Devraj, Rajesh, Ballwin, MO, UNITED STATES
 Flynn, Daniel L., Natick, MA, UNITED STATES
 Geng, Lifeng, Skokie, IL, UNITED STATES
 Graneto, Matthew J., Chesterfield, MO, UNITED STATES
 Hanau, Cathleen E., Chesterfield, MO, UNITED STATES
 Hanson, Gunnar J., Skokie, IL, UNITED STATES
 Hartmann, Susan J., Kirkwood, MO, UNITED STATES
 Hepperle, Michael, Boston, MA, UNITED STATES
 Huang, He, Chicago, IL, UNITED STATES
 Khanna, Ish K., Libertyville, IL, UNITED STATES
 Koszyk, Francis J., Prospect Heights, IL, UNITED STATES
 Liao, Shuyuan, Northbrook, IL, UNITED STATES
 Metz, Suzanne, Chesterfield, MO, UNITED STATES
 Naing, Win, Chesterfield, MO, UNITED STATES
 Partis, Richard A., Evanston, IL, UNITED STATES
 Perry, Thao D., Chesterfield, MO, UNITED STATES
 Rao, Shashidhar N., St. Louis, MO, UNITED STATES

Selness, Shaun Raj, Chesterfield, MO, UNITED STATES
South, Michael S., St. Louis, MO, UNITED STATES
Stealey, Michael A., Libertyville, IL, UNITED STATES
Talley, John Jeffrey, Cambridge, MA, UNITED STATES
Vazquez, Michael L., Ballwin, MO, UNITED STATES
Walker, John, Maryland Heights, MO, UNITED STATES
Weier, Richard M., Lake Bluff, IL, UNITED STATES
Xu, Xiangdong, Gurnee, IL, UNITED STATES
Yang, Syaulan, Chesterfield, MO, UNITED STATES
Yu, Yi, Glenville, IL, UNITED STATES
PATENT ASSIGNEE(S): Pharmacia Corporation, St. Louis, MO, UNITED STATES
(U.S. corporation)

| | NUMBER | KIND | DATE |
|-----------------------|--|------|---------------|
| PATENT INFORMATION: | US 7071198 | B1 | 20060704 |
| APPLICATION INFO.: | US 2004-840734 | | 20040505 (10) |
| RELATED APPLN. INFO.: | Division of Ser. No. US 2001-21780, filed on 7 Dec 2001, PENDING Division of Ser. No. US 2000-513351, filed on 24 Feb 2000, Pat. No. US 6525059 Continuation-in-part of Ser. No. US 1998-196623, filed on 20 Nov 1998, Pat. No. US 6514977 | | |

| | NUMBER | DATE |
|-----------------------|--|---------------|
| PRIORITY INFORMATION: | US 1997-47570P | 19970522 (60) |
| DOCUMENT TYPE: | Utility | |
| FILE SEGMENT: | GRANTED | |
| PRIMARY EXAMINER: | Solola, Taofiq | |
| LEGAL REPRESENTATIVE: | Lappin, Julie M. | |
| NUMBER OF CLAIMS: | 7 | |
| EXEMPLARY CLAIM: | 1 | |
| NUMBER OF DRAWINGS: | 0 Drawing Figure(s); 0 Drawing Page(s) | |
| LINE COUNT: | 18479 | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB A class of pyrazole derivatives is described for use in treating p38 kinase mediated disorders. Compounds of particular interest are defined by Formula IA ##STR1## wherein R.sup.1, R.sup.2, R.sup.3 and R.sup.4 are as described in the specification.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 4 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2005:197195 USPATFULL

TITLE: Mammalian tribbles signaling pathways and methods and reagents related thereto

INVENTOR(S): Dower, Steven, Sheffield, UNITED KINGDOM
Quanstrom, Eva, Sheffield, UNITED KINGDOM
Kiss-Toth, Endre, Sheffield, UNITED KINGDOM

| | NUMBER | KIND | DATE |
|---------------------|----------------|------|---------------|
| PATENT INFORMATION: | US 2005171338 | A1 | 20050804 |
| APPLICATION INFO.: | US 2003-466020 | A1 | 20020108 (10) |
| | WO 2002-US70 | | 20020108 |

| | NUMBER | DATE |
|-----------------------|-----------------|---------------|
| PRIORITY INFORMATION: | US 2001-260294P | 20010108 (60) |
| DOCUMENT TYPE: | Utility | |

<--

FILE SEGMENT: APPLICATION
LEGAL REPRESENTATIVE: MINTZ, LEVIN, COHN, FERRIS, GLOVSKY, AND POPEO, P.C.,
ONE FINANCIAL CENTER, BOSTON, MA, 02111, US
NUMBER OF CLAIMS: 41
EXEMPLARY CLAIM: 1
NUMBER OF DRAWINGS: 12 Drawing Page(s)
LINE COUNT: 5751

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB The invention provides methods and reagents for modulating mitogen
activated protein kinase pathways using mammalian tribbles homologs
(htrb).

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 5 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2005:171875 USPATFULL
TITLE: Jnk inhibitor
INVENTOR(S): Itoh, Fumio, Ibaraki, JAPAN
Kimura, Hiroyuki, Osaka, JAPAN
Igata, Hideki, Osaka, JAPAN
Kawamoto, Tomohiro, Osaka, JAPAN
Sasaki, Mitsuru, Osaka, JAPAN
Kitamura, Shuji, Osaka, JAPAN

| | NUMBER | KIND | DATE |
|---------------------|----------------|------|---------------|
| PATENT INFORMATION: | US 2005148624 | A1 | 20050707 |
| APPLICATION INFO.: | US 2003-504132 | A1 | 20030212 (10) |
| | WO 2003-JP1429 | | 20030212 |

| | NUMBER | DATE |
|-----------------------|--|----------|
| PRIORITY INFORMATION: | JP 2002-35073 | 20020213 |
| | JP 2003-2002251997 | 20020829 |
| DOCUMENT TYPE: | Utility | |
| FILE SEGMENT: | APPLICATION | |
| LEGAL REPRESENTATIVE: | TAKEDA PHARMACEUTICALS NORTH AMERICA, INC, INTELLECTUAL PROPERTY DEPARTMENT, 475 HALF DAY ROAD, SUITE 500, LINCOLNSHIRE, IL, 60069, US | |
| NUMBER OF CLAIMS: | 72 | |
| EXEMPLARY CLAIM: | 1 | |
| LINE COUNT: | 11597 | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB A JNK inhibitor containing a compound having an isoquinolinone skeleton
or a salt thereof, such as a compound represented by the formula
##STR1## wherein ring A and ring B are each an optionally substituted
benzene ring, X is --O--, --N.dbd., --NR.sup.3-- or --CHR.sup.3--,
R.sup.2 is an acyl group, an optionally esterified or thioesterified
carboxyl group, an optionally substituted carbamoyl group or an
optionally substituted amino group and the like, a broken line shows a
single bond or a double bond, and R.sup.1 is a hydrogen atom, an
optionally substituted hydrocarbon group, an optionally substituted
heterocyclic group and the like, and the like.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 6 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2005:171787 USPATFULL
TITLE: Methods for treating and preventing insulin resistance
and related disorders

INVENTOR(S): Greenberg, Andrew S., Boston, MA, UNITED STATES

| | NUMBER | KIND | DATE |
|-----------------------|---|------|---------------|
| PATENT INFORMATION: | US 2005148536 | A1 | 20050707 |
| APPLICATION INFO.: | US 2004-977116 | A1 | 20041029 (10) |
| RELATED APPLN. INFO.: | Continuation of Ser. No. US 2000-690647, filed on 17 Oct 2000, PENDING Continuation-in-part of Ser. No. WO 1999-US8364, filed on 16 Apr 1999, PENDING | | |

| | NUMBER | DATE | |
|-----------------------|---|---------------|-----|
| PRIORITY INFORMATION: | US 1998-82152P | 19980417 (60) | <-- |
| | US 1998-82741P | 19980423 (60) | <-- |
| DOCUMENT TYPE: | Utility | | |
| FILE SEGMENT: | APPLICATION | | |
| LEGAL REPRESENTATIVE: | FOLEY HOAG, LLP, PATENT GROUP, WORLD TRADE CENTER WEST, 155 SEAPORT BLVD, BOSTON, MA, 02110, US | | |
| NUMBER OF CLAIMS: | 18 | | |
| EXEMPLARY CLAIM: | 1 | | |
| NUMBER OF DRAWINGS: | 14 Drawing Page(s) | | |
| LINE COUNT: | 2690 | | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB The invention provides methods, therapeutics and kits for treating and preventing diseases or conditions associated with excessive lipolysis, in particular TNF- α induced lipolysis, and/or excessive free fatty acid levels. Exemplary conditions include insulin-resistance, **diabetes**, in particular NIDDM, obesity, glucose intolerance, hyperinsulinemia, polycystic ovary syndrome, and coronary artery disease. In a preferred embodiment, the method includes administering to a subject in need a pharmaceutically effective amount of an inhibitor of the JNK signal transduction pathway and/or an inhibitor of the MAPK/ERK signal transduction pathway.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 7 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2005:43617 USPATFULL

TITLE: Nucleic acid array

INVENTOR(S): Stuhlmuller, Bruno, Berlin, GERMANY, FEDERAL REPUBLIC OF

Haupl, Thomas, Erkner, GERMANY, FEDERAL REPUBLIC OF
PATENT ASSIGNEE(S): PathoArray GmbH, Berlin, GERMANY, FEDERAL REPUBLIC OF (non-U.S. corporation)

| | NUMBER | KIND | DATE |
|---------------------|----------------|------|---------------|
| PATENT INFORMATION: | US 2005037344 | A1 | 20050217 |
| APPLICATION INFO.: | US 2002-278698 | A1 | 20021023 (10) |

| | NUMBER | DATE | |
|-----------------------|--|----------|-----|
| PRIORITY INFORMATION: | DE 2001-10155600 | 20011109 | <-- |
| DOCUMENT TYPE: | Utility | | |
| FILE SEGMENT: | APPLICATION | | |
| LEGAL REPRESENTATIVE: | WELSH & KATZ, LTD, 120 S RIVERSIDE PLAZA, 22ND FLOOR, CHICAGO, IL, 60606 | | |
| NUMBER OF CLAIMS: | 29 | | |
| EXEMPLARY CLAIM: | 1 | | |
| LINE COUNT: | 2642 | | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB It is proposed to use selected monocyte macrophage genes to provide tools for diagnostic, prognostic and therapy-monitoring analysis and for performing screenings for pharmacologically active substances and substance classes of chronic inflammatory diseases, chronic inflammatory diseases induced by bacteria, arteriosclerosis, tumors, organ and tissue transplantations, and sepsis when examining blood, tissue, purified or cultivated cells.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 8 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2005:126974 USPATFULL

TITLE: Methods for treating and preventing insulin resistance and related disorders

INVENTOR(S): Greenberg, Andrew S., 711 Washington St., Boston, MA, UNITED STATES 02111

| | NUMBER | KIND | DATE |
|-----------------------|--|------|--------------|
| PATENT INFORMATION: | US 6897019 | B1 | 20050524 |
| APPLICATION INFO.: | US 2000-690647 | | 20001017 (9) |
| RELATED APPLN. INFO.: | Continuation-in-part of Ser. No. WO 1999-US8364, filed on 16 Apr 1999, PENDING | | |

| | NUMBER | DATE | |
|-----------------------|--|---------------|-----|
| PRIORITY INFORMATION: | US 1998-82741P | 19980423 (60) | <-- |
| | US 1998-82152P | 19980417 (60) | <-- |
| DOCUMENT TYPE: | Utility | | |
| FILE SEGMENT: | GRANTED | | |
| PRIMARY EXAMINER: | LeGuyader, John L. | | |
| ASSISTANT EXAMINER: | Schultz, J D | | |
| NUMBER OF CLAIMS: | 4 | | |
| EXEMPLARY CLAIM: | 1 | | |
| NUMBER OF DRAWINGS: | 26 Drawing Figure(s); 14 Drawing Page(s) | | |
| LINE COUNT: | 2976 | | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB The invention provides methods, therapeutics and kits for treating and preventing diseases or conditions associated with excessive lipolysis, in particular TNF- α induced lipolysis, and/or excessive free fatty acid levels. Exemplary conditions include insulin-resistance, **diabetes**, in particular NIDDM, obesity, glucose intolerance, hyperinsulinemia, polycystic ovary syndrome, and coronary artery disease. In a preferred embodiment, the method includes administering to a subject in need a pharmaceutically effective amount of an inhibitor of the JNK signal transduction pathway and/or an inhibitor of the MAPK/ERK signal transduction pathway.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 9 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2004:334786 USPATFULL

TITLE: Method for the detection of depression related gene transcripts in blood

INVENTOR(S): Liew, Choong-Chin, Toronto, CANADA

PATENT ASSIGNEE(S): ChondroGene Limited (non-U.S. corporation)

| NUMBER | KIND | DATE |
|--------|------|------|
| ----- | | |

PATENT INFORMATION: US 2004265868 A1 20041230
 APPLICATION INFO.: US 2004-812702 A1 20040330 (10)
 RELATED APPLN. INFO.: Continuation-in-part of Ser. No. US 2004-802875, filed
 on 12 Mar 2004, PENDING Continuation-in-part of Ser.
 No. US 2003-601518, filed on 20 Jun 2003, PENDING
 Continuation-in-part of Ser. No. US 2002-268730, filed
 on 9 Oct 2002, PENDING Continuation of Ser. No. US
 2000-477148, filed on 4 Jan 2000, ABANDONED

| | NUMBER | DATE | |
|-----------------------|--|---------------|-----|
| PRIORITY INFORMATION: | US 1999-115125P | 19990106 (60) | <-- |
| DOCUMENT TYPE: | Utility | | |
| FILE SEGMENT: | APPLICATION | | |
| LEGAL REPRESENTATIVE: | PALMER & DODGE, LLP, KATHLEEN M. WILLIAMS, 111 HUNTINGTON AVENUE, BOSTON, MA, 02199 | | |
| NUMBER OF CLAIMS: | 48 | | |
| EXEMPLARY CLAIM: | 1 | | |
| NUMBER OF DRAWINGS: | 34 Drawing Page(s) | | |
| LINE COUNT: | 5805 | | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB The present invention is directed to detection and measurement of gene transcripts and their equivalent nucleic acid products in blood. Specifically provided is analysis performed on a drop of blood for detecting, diagnosing and monitoring diseases using gene-specific and/or tissue-specific primers. The present invention also describes methods by which delineation of the sequence and/or quantitation of the expression levels of disease-specific genes allows for an immediate and accurate diagnostic/prognostic test for disease or to assess the effect of a particular treatment regimen.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 10 OF 34 USPATFULL on STN
 ACCESSION NUMBER: 2004:299993 USPATFULL
 TITLE: Protein kinase inhibitors
 INVENTOR(S): Burns, Christopher John, Seddon, AUSTRALIA
 Bu, Xianying, Rosanna East, AUSTRALIA
 Wilks, Andrew Frederick, South Yarra, AUSTRALIA

| | NUMBER | KIND | DATE |
|---------------------|----------------|------|---------------|
| PATENT INFORMATION: | US 2004235862 | A1 | 20041125 |
| APPLICATION INFO.: | US 2003-469303 | A1 | 20031204 (10) |
| | WO 2003-AU629 | | 20030523 |

| | NUMBER | DATE | |
|-----------------------|--|---------------|-----|
| PRIORITY INFORMATION: | AU 2002-2515 | 20020523 | <-- |
| | US 2002-399070P | 20020726 (60) | <-- |
| DOCUMENT TYPE: | Utility | | |
| FILE SEGMENT: | APPLICATION | | |
| LEGAL REPRESENTATIVE: | MORRISON & FOERSTER LLP, 3811 VALLEY CENTRE DRIVE, SUITE 500, SAN DIEGO, CA, 92130-2332 | | |
| NUMBER OF CLAIMS: | 25 | | |
| EXEMPLARY CLAIM: | CLM-01-24 | | |
| NUMBER OF DRAWINGS: | 3 Drawing Page(s) | | |
| LINE COUNT: | 1645 | | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB A compound of the general formula: ##STR1##

or pharmaceutically acceptable salts, hydrates, solvates, crystal forms of diastereomers thereof is described. A method of treating protein kinase-associated disease states using the compound of Formula I is also described.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 11 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2004:228047 USPATFULL

TITLE: Substituted pyrazoles as p38 kinase inhibitors

INVENTOR(S): Naraian, Ashok S., Ballwin, MO, UNITED STATES

Clare, Michael, Skokie, IL, UNITED STATES

Collins, Paul W., Deerfield, IL, UNITED STATES

Crich, Joyce Zuowu, Glenview, IL, UNITED STATES

Devraj, Rajesh, Ballwin, MO, UNITED STATES

Flynn, Daniel L., Natick, MA, UNITED STATES

Geng, Lifeng, Skokie, IL, UNITED STATES

Graneto, Matthew J., Chesterfield, MO, UNITED STATES

Hanau, Cathleen E., Chesterfield, MO, UNITED STATES

Hanson, Gunnar J., Skokie, IL, UNITED STATES

Hartmann, Susan J., Kirkwood, MO, UNITED STATES

Hepperle, Michael, Boston, MA, UNITED STATES

Huang, He, Northbrook, IL, UNITED STATES

Koszyk, Francis J., Prospect Heights, IL, UNITED STATES

Liao, Shuyuan, Northbrook, IL, UNITED STATES

Metz, Suzanne, Chesterfield, MO, UNITED STATES

Partis, Richard A., Evanston, IL, UNITED STATES

Perry, Thao D., Chesterfield, MO, UNITED STATES

Rao, Shashidhar N., St. Louis, MO, UNITED STATES

Selness, Shaun Raj, Chesterfield, MO, UNITED STATES

South, Michael S., St. Louis, MO, UNITED STATES

Stealey, Michael A., Libertyville, IL, UNITED STATES

Talley, John Jeffrey, Cambridge, MA, UNITED STATES

Vazquez, Michael L., Ballwin, MO, UNITED STATES

Weier, Richard M., Lake Bluff, IL, UNITED STATES

Xu, Xiangdong, Gurnee, IL, UNITED STATES

Khanna, Ish K., Libertyville, IL, UNITED STATES

Yu, Yi, Glenview, IL, UNITED STATES

Naing, Win, Chesterfield, MO, UNITED STATES

Yang, Syaulan, Chesterfield, MO, UNITED STATES

| | NUMBER | KIND | DATE |
|-----------------------|--|------|---------------|
| PATENT INFORMATION: | US 2004176433 | A1 | 20040909 |
| APPLICATION INFO.: | US 2003-374781 | A1 | 20030225 (10) |
| RELATED APPLN. INFO.: | Division of Ser. No. US 2002-114297, filed on 2 Apr 2002, GRANTED, Pat. No. US 6617324 Division of Ser. No. US 2001-918481, filed on 31 Jul 2001, GRANTED, Pat. No. US 6423713 Division of Ser. No. US 1998-196623, filed on 20 Nov 1998, GRANTED, Pat. No. US 6514977 Continuation-in-part of Ser. No. US 1998-83670, filed on 22 May 1998, ABANDONED | | |

| | NUMBER | DATE |
|-----------------------|---|---------------|
| PRIORITY INFORMATION: | US 1997-47570P | 19970522 (60) |
| DOCUMENT TYPE: | Utility | |
| FILE SEGMENT: | APPLICATION | |
| LEGAL REPRESENTATIVE: | David M. Gryte, Harness, Dickey & Pierce, P.L.C., Suite | |

400, 7700 Bonhomme, St. Louis, MO, 63105
NUMBER OF CLAIMS: 184
EXEMPLARY CLAIM: 1
NUMBER OF DRAWINGS: 2 Drawing Page(s)
LINE COUNT: 21540

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB A class of pyrazole derivatives is described for use in treating p38 kinase mediated disorders. Compounds of particular interest are defined by Formula IA ##STR1##

wherein R.sup.1, R.sup.2, R.sup.3 and R.sup.4 are as described in the specification.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 12 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2004:204017 USPATFULL
TITLE: 4-Aryl substituted indolinones
INVENTOR(S): Cui, Jingrong, Foster City, CA, UNITED STATES
Zhang, Ruofei, Foster City, CA, UNITED STATES
Shen, Hong, San Francisco, CA, UNITED STATES
Chu, Ji Yu, Fremont, CA, UNITED STATES
Zhang, Fang-Jie, San Jose, CA, UNITED STATES
Koenig, Marcel, Burlingame, CA, UNITED STATES
Do, Steven Huy, San Jose, CA, UNITED STATES
Li, Xiaoyuan, Los Altos, CA, UNITED STATES
Wei, Chung Chen, Foster City, CA, UNITED STATES
Tang, Peng Cho, Moraga, CA, UNITED STATES
PATENT ASSIGNEE(S): Sugan, Inc. (U.S. corporation)

| | NUMBER | KIND | DATE |
|-----------------------|--|------|---------------|
| PATENT INFORMATION: | US 2004157909 | A1 | 20040812 |
| | US 6861418 | B2 | 20050301 |
| APPLICATION INFO.: | US 2003-736243 | A1 | 20031216 (10) |
| RELATED APPLN. INFO.: | Division of Ser. No. US 2001-23488, filed on 20 Dec 2001, GRANTED, Pat. No. US 6677368 | | |

| | NUMBER | DATE |
|-----------------------|---|---------------|
| PRIORITY INFORMATION: | US 2000-256479P | 20001220 (60) |
| DOCUMENT TYPE: | Utility | |
| FILE SEGMENT: | APPLICATION | |
| LEGAL REPRESENTATIVE: | FOLEY AND LARDNER, SUITE 500, 3000 K STREET NW, WASHINGTON, DC, 20007 | |
| NUMBER OF CLAIMS: | 20 | |
| EXEMPLARY CLAIM: | 1 | |
| LINE COUNT: | 13339 | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB The present invention relates to 4-arylindolinones, as well as pharmaceutical compositions thereof, capable of modulating protein kinase signal transduction in order to regulate, modulate and/or inhibit abnormal cell proliferation. The present invention also relates to methods for treating protein kinase related disorders.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 13 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2004:77185 USPATFULL
TITLE: Substituted pyridinones

INVENTOR(S) : Devadas, Balekudru, Chesterfield, MO, UNITED STATES
Walker, John, Maryland Heights, MO, UNITED STATES
Selness, Shaun R., Chesterfield, MO, UNITED STATES
Boehm, Terri L., Ballwin, MO, UNITED STATES
Durley, Richard C., Chesterfield, MO, UNITED STATES
Devraj, Rajesh, Ballwin, MO, UNITED STATES
Hickory, Brian S., Wildwood, MO, UNITED STATES
Rucker, Paul V., University City, MO, UNITED STATES
Jerome, Kevin D., Maryland Heights, MO, UNITED STATES
Madsen, Heather M., University City, MO, UNITED STATES
Alvira, Edgardo, Chesterfield, MO, UNITED STATES
Promo, Michele A., Maryland Heights, MO, UNITED STATES
Blevis-Bal, Radhika M., St. Louis, MO, UNITED STATES
Marruto, Laura D., Ellisville, MO, UNITED STATES
Hitchcock, Jeff, Saint Peters, MO, UNITED STATES
Owen, Thomas, Chesterfield, MO, UNITED STATES
Naing, Win, Chesterfield, MO, UNITED STATES
Xing, Li, Chesterfield, MO, UNITED STATES
Shieh, Huey S., St. Louis, MO, UNITED STATES
Sambandam, Aruna, Guilderland, NY, UNITED STATES
Liu, Shuang, Schenectady, NY, UNITED STATES
Scott, Ian L., Woodinville, WA, UNITED STATES
McGee, Kevin F., Guilderland, NY, UNITED STATES

| | NUMBER | KIND | DATE |
|---------------------|----------------|------|---------------|
| PATENT INFORMATION: | US 2004058964 | A1 | 20040325 |
| | US 7067540 | B2 | 20060627 |
| APPLICATION INFO.: | US 2003-367987 | A1 | 20030214 (10) |

| | NUMBER | DATE |
|--|--|---------------|
| PRIORITY INFORMATION: | US 2002-357029P | 20020214 (60) |
| | US 2002-436915P | 20021230 (60) |
| DOCUMENT TYPE: | Utility | |
| FILE SEGMENT: | APPLICATION | |
| LEGAL REPRESENTATIVE: | Steven J. Sarussi, McDonnell Boehnen Hulbert & Berghoff, 32nd Floor, 300 S. Wacker Drive, Chicago, IL, 60606 | |
| NUMBER OF CLAIMS: | 76 | |
| EXEMPLARY CLAIM: | 1 | |
| LINE COUNT: | 26020 | |
| CAS INDEXING IS AVAILABLE FOR THIS PATENT. | | |
| AB | Disclosed are compounds Formula I | ##STR1## |

and pharmaceutically acceptable salts thereof, wherein R.sub.1, R.sub.2, R.sub.3, R.sub.4, and R.sub.5 are defined herein. These compounds are useful for treating diseases and conditions caused or exacerbated by unregulated p38 MAP Kinase and/or TNF activity. Pharmaceutical compositions containing the compounds, methods of preparing the compounds and methods of treatment using the compounds are also disclosed.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 14 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2003:300912 USPATFULL

TITLE: Combinations of peroxisome proliferator-activated
receptor-alpha agonists and cyclooxygenase-2 selective
inhibitors and therapeutic uses therefor

INVENTOR(S): Obukowicz, Mark G., Kirkwood, MO, UNITED STATES
PATENT ASSIGNEE(S): Pharmacia Corporation, St. Louis, MO, UNITED STATES,
63141 (U.S. corporation)

| | NUMBER | KIND | DATE |
|---------------------|----------------|------|---------------|
| PATENT INFORMATION: | US 2003212138 | A1 | 20031113 |
| APPLICATION INFO.: | US 2003-341217 | A1 | 20030113 (10) |

| | NUMBER | DATE |
|-----------------------|---|-------------------|
| PRIORITY INFORMATION: | US 2002-348297P | 20020114 (60) <-- |
| DOCUMENT TYPE: | Utility | |
| FILE SEGMENT: | APPLICATION | |
| LEGAL REPRESENTATIVE: | Charles E. Dunlap, Keenan Building, Third Floor, 1330 Lady Street, Columbia, SC, 29201 | |
| NUMBER OF CLAIMS: | 59 | |
| EXEMPLARY CLAIM: | 1 | |
| LINE COUNT: | 4257 | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB Methods for the treatment, prevention, or inhibition of pain, inflammation, or inflammation-related disorder, and for the treatment or inhibition of cardiovascular disease or disorder, and for the treatment or inhibition of cancer, and for the treatment of Alzheimer's disease in a subject in need of such treatment, prevention, or inhibition, include treating the subject with a peroxisome proliferator activated receptor- α agonist and a cyclooxygenase-2 selective inhibitor or prodrug thereof. Compositions, pharmaceutical compositions and kits for effecting the particular methods are also described.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 15 OF 34 USPATFULL on STN
ACCESSION NUMBER: 2003:225250 USPATFULL
TITLE: Combinations of a cyclooxygenase-2 selective inhibitor and a TNF α antagonist and therapeutic uses therefor
INVENTOR(S): Bennett, Dennis A., Wildwood, MO, UNITED STATES
PATENT ASSIGNEE(S): Pharmacia Corporation, St. Louis, MO, UNITED STATES
(U.S. corporation)

| | NUMBER | KIND | DATE |
|---------------------|----------------|------|---------------|
| PATENT INFORMATION: | US 2003157061 | A1 | 20030821 |
| APPLICATION INFO.: | US 2002-310454 | A1 | 20021205 (10) |

| | NUMBER | DATE |
|-----------------------|---|-------------------|
| PRIORITY INFORMATION: | US 2001-337802P | 20011205 (60) <-- |
| DOCUMENT TYPE: | Utility | |
| FILE SEGMENT: | APPLICATION | |
| LEGAL REPRESENTATIVE: | Charles E. Dunlap, Keenan Building, Third Floor, 1330 Lady Street, Columbia, SC, 29201 | |
| NUMBER OF CLAIMS: | 75 | |
| EXEMPLARY CLAIM: | 1 | |
| LINE COUNT: | 3289 | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB A method for the prevention, treatment, or inhibition of pain, inflammation, or inflammation-related disorder and for the prevention, treatment, or inhibition of a cardiovascular disease or disorder in a subject that is in need of such prevention, treatment or inhibition,

involves the administration to the subject of a cyclooxygenase-2 selective inhibitor or prodrug thereof and a TNF α antagonist. A method can also involve the treatment, prevention, or inhibition of cancer in a subject in need of such treatment, prevention, or inhibition, by administering to the subject a cyclooxygenase-2 selective inhibitor or prodrug thereof and a TNF α antagonist which is selected from the group consisting of a compound that affects the synthesis of TNF α , a compound that inhibits the binding of TNF α with a receptor specific for TNF α , and a compound that interferes with intracellular signaling triggered by TNF α binding with a receptor. Compositions, pharmaceutical compositions and kits that can be used with the methods are also described.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 16 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2003:187403 USPATFULL
 TITLE: Tumor necrosis factor-gamma
 INVENTOR(S): Yu, Guo-Liang, Berkeley, CA, UNITED STATES
 Ni, Jian, Germantown, MD, UNITED STATES
 Rosen, Craig A., Laytonsville, MD, UNITED STATES
 Zhang, Jun, San Diego, CA, UNITED STATES

| | NUMBER | KIND | DATE |
|-----------------------|---|------|---------------|
| PATENT INFORMATION: | US 2003129189 | A1 | 20030710 |
| APPLICATION INFO.: | US 2002-226294 | A1 | 20020823 (10) |
| RELATED APPLN. INFO.: | Continuation-in-part of Ser. No. US 2001-899059, filed on 6 Jul 2001, PENDING Continuation-in-part of Ser. No. US 2000-559290, filed on 27 Apr 2000, ABANDONED Continuation-in-part of Ser. No. US 1999-246129, filed on 8 Feb 1999, PENDING Continuation-in-part of Ser. No. US 1998-131237, filed on 7 Aug 1998, PENDING Continuation-in-part of Ser. No. US 1998-5020, filed on 9 Jan 1998, ABANDONED Continuation-in-part of Ser. No. US 1995-461246, filed on 5 Jun 1995, ABANDONED Continuation-in-part of Ser. No. WO 1994-US12880, filed on 7 Nov 1994, PENDING | | |

| | NUMBER | DATE | |
|-----------------------|---|----------|----------|
| PRIORITY INFORMATION: | US 2001-314381P | 20010824 | (60) <-- |
| | US 2001-278449P | 20010326 | (60) <-- |
| | US 2000-216879P | 20000707 | (60) <-- |
| | US 2000-180908P | 20000208 | (60) <-- |
| | US 1999-134067P | 19990513 | (60) <-- |
| | US 1999-132227P | 19990503 | (60) <-- |
| | US 1999-131963P | 19990430 | (60) <-- |
| | US 1998-74047P | 19980209 | (60) <-- |
| DOCUMENT TYPE: | Utility | | |
| FILE SEGMENT: | APPLICATION | | |
| LEGAL REPRESENTATIVE: | HUMAN GENOME SCIENCES INC, 9410 KEY WEST AVENUE, ROCKVILLE, MD, 20850 | | |
| NUMBER OF CLAIMS: | 49 | | |
| EXEMPLARY CLAIM: | 1 | | |
| NUMBER OF DRAWINGS: | 33 Drawing Page(s) | | |
| LINE COUNT: | 13325 | | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB Human TNF-gamma-alpha and TNF-gamma-beta polypeptides and DNA (RNA) encoding such polypeptides and a procedure for producing such

polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing such polypeptides to inhibit cellular growth, for example in a tumor or cancer, for facilitating wound-healing, to provide resistance against infection, induce inflammatory activities, and stimulating the growth of certain cell types to treat diseases, for example restenosis. Also disclosed are diagnostic methods for detecting a mutation in the TNF-gamma-alpha and TNF-gamma-beta nucleic acid sequences or overexpression of the TNF-gamma-alpha and/or TNF-gamma-beta polypeptides. Antagonists against such polypeptides and their use as a therapeutic to treat cachexia, septic shock, cerebral malaria, inflammation, arthritis and graft-rejection are also disclosed.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 17 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2003:100186 USPATFULL
TITLE: 4-aryl substituted indolinones
INVENTOR(S): Cui, Jingrong Jean, Foster City, CA, UNITED STATES
Zhang, Ruofei, Foster City, CA, UNITED STATES
Shen, Hong, San Francisco, CA, UNITED STATES
Chu, Ji Yu, Fremont, CA, UNITED STATES
Zhang, Fang-Jie, San Jose, CA, UNITED STATES
Koenig, Marcel, Burlingame, CA, UNITED STATES
Do, Steven Huy, San Jose, CA, UNITED STATES
Li, Xiaoyuan, Los Altos, CA, UNITED STATES
Wei, Chung Chen, Foster City, CA, UNITED STATES
Tang, Peng Cho, Moraga, CA, UNITED STATES
PATENT ASSIGNEE(S): Sugan, Inc. (U.S. corporation)

| | NUMBER | KIND | DATE |
|---------------------|---------------|------|---------------|
| PATENT INFORMATION: | US 2003069297 | A1 | 20030410 |
| | US 6677368 | B2 | 20040113 |
| APPLICATION INFO.: | US 2001-23488 | A1 | 20011220 (10) |

| | NUMBER | DATE |
|-----------------------|--|---------------|
| PRIORITY INFORMATION: | US 2000-256479P | 20001220 (60) |
| DOCUMENT TYPE: | Utility | |
| FILE SEGMENT: | APPLICATION | |
| LEGAL REPRESENTATIVE: | FOLEY AND LARDNER, SUITE 500, 3000 K STREET NW, WASHINGTON, DC, 20007 | |
| NUMBER OF CLAIMS: | 23 | |
| EXEMPLARY CLAIM: | 1 | |
| LINE COUNT: | 14189 | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB The present invention relates to 4-arylindolinones, as well as pharmaceutical compositions thereof, capable of modulating protein kinase signal transduction in order to regulate, modulate and/or inhibit abnormal cell proliferation. The present invention also relates to methods for treating protein kinase related disorders.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 18 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2003:79103 USPATFULL
TITLE: Use of carboxy compounds such as 2(4-acetoxyphenyl)-2-chloro-N-methyl-ethylammonium chloride as anti-inflammatory agents
INVENTOR(S): De Bosscher, Karolien, Zottegem, BELGIUM

Berghe, Wim Vanden, Gentbrugge, BELGIUM
 Haegeman, Guy, Balegem, BELGIUM

| | NUMBER | KIND | DATE |
|-----------------------|---|------|---------------|
| PATENT INFORMATION: | US 2003055030 | A1 | 20030320 |
| | US 7053120 | B2 | 20060530 |
| APPLICATION INFO.: | US 2002-177987 | A1 | 20020621 (10) |
| RELATED APPLN. INFO.: | Continuation of Ser. No. WO 2000-EP13347, filed on 21 Dec 2000, UNKNOWN | | |

| | NUMBER | DATE | |
|-----------------------|--|----------|-----|
| PRIORITY INFORMATION: | EP 1999-204433 | 19991221 | <-- |
| DOCUMENT TYPE: | Utility | | |
| FILE SEGMENT: | APPLICATION | | |
| LEGAL REPRESENTATIVE: | TRASKBRITT, PC, P.O. Box 2550, Salt Lake City, UT, 84110 | | |
| NUMBER OF CLAIMS: | 8 | | |
| EXEMPLARY CLAIM: | 1 | | |
| NUMBER OF DRAWINGS: | 6 Drawing Page(s) | | |
| LINE COUNT: | 1147 | | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB The present invention relates to the use of specific carboxy compounds, more specifically to the use of 2(4-acetoxyphenyl)-2-chloro-N-methyl-ethylammonium chloride, in the treatment of inflammatory diseases. Part of the invention is also a composition, preferably a pharmaceutical composition, comprising as active ingredient at least 2 (4-acetoxyphenyl)-2-chloro-N-methyl-ethylammonium chloride together with (pharmaceutically) acceptable excipients.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 19 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2003:57074 USPATFULL
 TITLE: Carbon monoxide improves outcomes in tissue and organ transplants and suppresses apoptosis
 INVENTOR(S): Bach, Fritz H., Manchester-by-the-Sea, MA, UNITED STATES
 Otterbein, Leo E., New Kensington, PA, UNITED STATES
 Soares, Miguel P., Boston, MA, UNITED STATES
 Tobiasch, Edda M., Bonn, GERMANY, FEDERAL REPUBLIC OF
 Gose, Jeanne, Manchester-by-the-Sea, MA, UNITED STATES

| | NUMBER | KIND | DATE |
|---------------------|----------------|------|---------------|
| PATENT INFORMATION: | US 2003039638 | A1 | 20030227 |
| APPLICATION INFO.: | US 2002-177930 | A1 | 20020621 (10) |

| | NUMBER | DATE | |
|-----------------------|--|---------------|-----|
| PRIORITY INFORMATION: | US 2001-300289P | 20010621 (60) | <-- |
| | US 2001-334340P | 20011129 (60) | <-- |
| | US 2001-337974P | 20011207 (60) | <-- |
| DOCUMENT TYPE: | Utility | | |
| FILE SEGMENT: | APPLICATION | | |
| LEGAL REPRESENTATIVE: | FISH & RICHARDSON PC, 225 FRANKLIN ST, BOSTON, MA, 02110 | | |
| NUMBER OF CLAIMS: | 149 | | |
| EXEMPLARY CLAIM: | 1 | | |

NUMBER OF DRAWINGS: 31 Drawing Page(s)

LINE COUNT: 3473

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB The present invention features methods for transplanting organs, tissues and individual cells. Also featured are methods for maintaining cells in vitro and for enhancing survival and/or function of cells following transplantation. The methods include the administration of carbon monoxide in an amount sufficient to enhance cell survival and/or function.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 20 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2003:51146 USPATFULL

TITLE: Methods and compositions for diagnosis and treatment of vascular conditions

INVENTOR(S): Pillarisetti, Sivaram, Norcross, GA, UNITED STATES
Wang, Dongyan, Norcross, GA, UNITED STATES
Saxena, Uday, Atlanta, GA, UNITED STATES

| | NUMBER | KIND | DATE |
|---------------------|----------------|------|---------------|
| PATENT INFORMATION: | US 2003036103 | A1 | 20030220 |
| APPLICATION INFO.: | US 2002-210896 | A1 | 20020731 (10) |

| | NUMBER | DATE |
|-----------------------|---|-------------------|
| PRIORITY INFORMATION: | US 2001-309012P | 20010731 (60) <-- |
| DOCUMENT TYPE: | Utility | |
| FILE SEGMENT: | APPLICATION | |
| LEGAL REPRESENTATIVE: | JOHN S. PRATT, ESQ, KILPATRICK STOCKTON, LLP, 1100 PEACHTREE STREET, SUITE 2800, ATLANTA, GA, 30309 | |
| NUMBER OF CLAIMS: | 18 | |
| EXEMPLARY CLAIM: | 1 | |
| NUMBER OF DRAWINGS: | 6 Drawing Page(s) | |
| LINE COUNT: | 1032 | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB The present invention is directed to methods and compositions for the diagnosis and treatment of vascular conditions, particularly **diabetes** and atherosclerosis. The present invention comprises methods and compositions for determining the expression or activity of enzymes effecting HSPG, preferably, heparanase. The invention also comprises methods and compositions for treatment of vasculophathic diseases comprising administration of therapeutic compounds that are effective in inhibiting the expression or activity of heparanase.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 21 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2003:37639 USPATFULL

TITLE: TUMOR NECROSIS FACTOR-GAMMA

INVENTOR(S): YU, GUO-LIANG, SAN MATEO, CA, UNITED STATES
NI, JIAN, ROCKVILLE, MD, UNITED STATES
ROSEN, CRAIG A., LAYTONSVILLE, MD, UNITED STATES
ZHANG, JUN, BETHESDA, MD, UNITED STATES

| | NUMBER | KIND | DATE |
|---------------------|---------------|------|----------|
| PATENT INFORMATION: | US 2003027284 | A1 | 20030206 |
| | US 6599719 | B2 | 20030729 |

APPLICATION INFO.: US 1998-131237 A1 19980807 (9)
 RELATED APPLN. INFO.: Continuation-in-part of Ser. No. US 1998-5020, filed on
 9 Jan 1998, ABANDONED Continuation-in-part of Ser. No.
 US 1995-461246, filed on 5 Jun 1995, ABANDONED
 Continuation-in-part of Ser. No. WO 1994-US12880, filed
 on 7 Nov 1994, UNKNOWN

| | NUMBER | DATE | |
|-----------------------|--|---------------|-----|
| | ----- | ----- | |
| PRIORITY INFORMATION: | US 1998-74047P | 19980209 (60) | <-- |
| DOCUMENT TYPE: | Utility | | |
| FILE SEGMENT: | APPLICATION | | |
| LEGAL REPRESENTATIVE: | HUMAN GENOME SCIENCES INC, 9410 KEY WEST AVENUE, ROCKVILLE, MD, 20850 | | |
| NUMBER OF CLAIMS: | 41 | | |
| EXEMPLARY CLAIM: | 1 | | |
| NUMBER OF DRAWINGS: | 30 Drawing Page(s) | | |
| LINE COUNT: | 6325 | | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB Human TNF-gamma-alpha and TNF-gamma-beta polypeptides and DNA (RNA) encoding such polypeptides and a procedure for producing such polypeptides by recombinant techniques is disclosed. Also disclosed are methods for utilizing such polypeptides to inhibit cellular growth, for example in a tumor or cancer, for facilitating wound-healing, to provide resistance against infection, induce inflammatory activities, and stimulating the growth of certain cell types to treat diseases, for example restenosis. Also disclosed are diagnostic methods for detecting a mutation in the TNF-gamma-alpha and TNF-gamma-beta nucleic acid sequences or overexpression of the TNF-gamma-alpha and TNF-gamma-beta polypeptides. Antagonists against such polypeptides and their use as a therapeutic to treat cachexia, septic shock, cerebral malaria, inflammation, arthritis and graft-rejection are also disclosed.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 22 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2003:240399 USPATFULL

TITLE: Substituted pyrazoles as p38 kinase inhibitors

INVENTOR(S): Naraian, Ashok S., Hainesville, IL, United States

Clare, Michael, Skokie, IL, United States

Collins, Paul W., Deerfield, IL, United States

Crich, Joyce Zuowu, Glenview, IL, United States

Devraj, Rajesh, Ballwin, MO, United States

Flynn, Daniel L., Clarkson Valley, MO, United States

Geng, Lifeng, Skokie, IL, United States

Graneto, Matthew J., Chesterfield, MO, United States

Hanau, Cathleen E., Chesterfield, MO, United States

Hanson, Gunnar J., Skokie, IL, United States

Hartmann, Susan J., Kirkwood, MO, United States

Hepperle, Michael, St. Charles, MO, United States

Huang, He, Chicago, IL, United States

Koszyk, Francis J., Prospect Heights, IL, United States

Liao, Shuyuan, Glen Ellyn, IL, United States

Metz, Suzanne, Chesterfield, MO, United States

Partis, Richard A., Evanston, IL, United States

Perry, Thao D., Chesterfield, MO, United States

Rao, Shashidhar N., St. Louis, MO, United States

Selness, Shaun Raj, Chesterfield, MO, United States

South, Michael S., St. Louis, MO, United States

Stealey, Michael A., Libertyville, IL, United States

Talley, John Jeffrey, St. Louis, MO, United States
Vazquez, Michael L., Ballwin, MO, United States
Weier, Richard M., Lake Bluff, IL, United States
Xi, Xiangdong, Evanston, IL, United States
Khanna, Ish K., Libertyville, IL, United States
Yu, Yi, Skokie, IL, United States
PATENT ASSIGNEE(S): G. D. Searle & Company, Skokie, IL, United States (U.S. corporation)

| | NUMBER | KIND | DATE |
|-----------------------|--|------|---------------|
| PATENT INFORMATION: | US 6617324 | B1 | 20030909 |
| APPLICATION INFO.: | US 2002-114297 | | 20020402 (10) |
| RELATED APPLN. INFO.: | Division of Ser. No. US 2001-918481, filed on 31 Jul 2001, now patented, Pat. No. US 6423713 Division of Ser. No. US 1998-196623, filed on 20 Nov 1998 Continuation-in-part of Ser. No. US 1998-83670, filed on 22 May 1998, now abandoned | | |

| | NUMBER | DATE |
|--|--|---------------|
| PRIORITY INFORMATION: | US 1997-47570P | 19970522 (60) |
| DOCUMENT TYPE: | Utility | |
| FILE SEGMENT: | GRANTED | |
| PRIMARY EXAMINER: | Solola, T. A. | |
| LEGAL REPRESENTATIVE: | Gryte, David M., Harness, Dickey & Pierce, P.L.C. | |
| NUMBER OF CLAIMS: | 112 | |
| EXEMPLARY CLAIM: | 1 | |
| NUMBER OF DRAWINGS: | 2 Drawing Figure(s); 2 Drawing Page(s) | |
| LINE COUNT: | 17190 | |
| CAS INDEXING IS AVAILABLE FOR THIS PATENT. | | |
| AB | A class of pyrazole derivatives described for use in treating p38 kinase mediated disorders. Compounds of particular interest are defined by Formula IA ##STR1## | |

wherein R.sup.1, R.sup.2, R.sup.3 and R.sup.4 are as described in the specification.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 23 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2003:33482 USPATFULL
TITLE: Substituted pyrazoles as p38 kinase inhibitors
INVENTOR(S): Anantanarayan, Ashok, Hainesville, IL, United States
Clare, Michael, Skokie, IL, United States
Collins, Paul W., Deerfield, IL, United States
Crich, Joyce Zuowu, Glenview, IL, United States
Devraj, Rajesh, Ballwin, MO, United States
Flynn, Daniel L., Clarkson Valley, MO, United States
Geng, Lifeng, Skokie, IL, United States
Graneto, Matthew J., Chesterfield, MO, United States
Hanau, Cathleen E., Chesterfield, MO, United States
Hanson, Gunnar J., Skokie, IL, United States
Hartmann, Susan J., Kirkwood, MO, United States
Hepperle, Michael, St. Charles, MO, United States
Huang, He, Chicago, IL, United States
Koszyk, Francis J., Prospect Heights, IL, United States
Liao, Shuyuan, Glen Ellyn, IL, United States
Metz, Suzanne, Chesterfield, MO, United States
Partis, Richard A., Evanston, IL, United States

Perry, Thao D., Chesterfield, MO, United States
 Rao, Shashidhar N., St. Louis, MO, United States
 Selness, Shaun Raj, Chesterfield, MO, United States
 South, Michael S., St. Louis, MO, United States
 Stealey, Michael A., Libertyville, IL, United States
 Talley, John Jeffrey, St. Louis, MO, United States
 Vazquez, Michael L., Ballwin, MO, United States
 Weier, Richard M., Lake Bluff, IL, United States
 Xu, Xiangdong, Evanston, IL, United States
 Khanna, Ish K., Libertyville, IL, United States
 Yu, Yi, Skokie, IL, United States
 PATENT ASSIGNEE(S): G.D. Searle & Company, Skokie, IL, United States (U.S. corporation)

| | NUMBER | KIND | DATE |
|-----------------------|--|------|--------------|
| PATENT INFORMATION: | US 6514977 | B1 | 20030204 |
| APPLICATION INFO.: | US 1998-196623 | | 19981120 (9) |
| RELATED APPLN. INFO.: | Continuation-in-part of Ser. No. US 1998-83670, filed on 22 May 1998 | | |

| | NUMBER | DATE |
|-----------------------|--|---------------|
| PRIORITY INFORMATION: | US 1997-47570P | 19970522 (60) |
| DOCUMENT TYPE: | Utility | |
| FILE SEGMENT: | GRANTED | |
| PRIMARY EXAMINER: | Solola, Taofiq A. | |
| LEGAL REPRESENTATIVE: | Williams, Scott A. | |
| NUMBER OF CLAIMS: | 92 | |
| EXEMPLARY CLAIM: | 1 | |
| NUMBER OF DRAWINGS: | 2 Drawing Figure(s); 2 Drawing Page(s) | |
| LINE COUNT: | 16885 | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB A class of pyrazole derivatives is described for use in treating p38 kinase mediated disorders. Compounds of particular interest are defined by Formula IA ##STR1##

wherein R.sup.1, R.sup.2, R.sup.3 and R.sup.4 are as described in the specification.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 24 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2002:272419 USPATFULL
 TITLE: Tumor necrosis factor-gamma
 INVENTOR(S): Yu, Guo-Liang, Berkeley, CA, UNITED STATES
 Ni, Jian, Germantown, MD, UNITED STATES
 Rosen, Craig A., Laytonsville, MD, UNITED STATES
 Zhang, Jun, Bethesda, MD, UNITED STATES

| | NUMBER | KIND | DATE |
|-----------------------|---|------|--------------|
| PATENT INFORMATION: | US 2002150534 | A1 | 20021017 |
| APPLICATION INFO.: | US 2001-899059 | A1 | 20010706 (9) |
| RELATED APPLN. INFO.: | Continuation-in-part of Ser. No. WO 2000-US11689, filed on 28 Apr 2000, UNKNOWN Continuation-in-part of Ser. No. US 1999-246129, filed on 8 Feb 1999, PENDING Continuation-in-part of Ser. No. US 1998-131237, filed on 7 Aug 1998, PENDING Continuation-in-part of Ser. No. US 1998-5020, filed on 9 Jan 1998, ABANDONED | | |

Continuation-in-part of Ser. No. US 1995-461246, filed
on 5 Jun 1995, ABANDONED Continuation-in-part of Ser.
No. WO 1994-US12880, filed on 7 Nov 1994, UNKNOWN

| | NUMBER | DATE | |
|-----------------------|--|---------------|-----|
| PRIORITY INFORMATION: | US 2001-278449P | 20010326 (60) | <-- |
| | US 2000-216879P | 20000707 (60) | <-- |
| | US 2000-180908P | 20000208 (60) | <-- |
| | US 1999-134067P | 19990513 (60) | <-- |
| | US 1999-132227P | 19990503 (60) | <-- |
| | US 1999-131963P | 19990430 (60) | <-- |
| | US 1998-74047P | 19980209 (60) | <-- |
| DOCUMENT TYPE: | Utility | | |
| FILE SEGMENT: | APPLICATION | | |
| LEGAL REPRESENTATIVE: | HUMAN GENOME SCIENCES INC, 9410 KEY WEST AVENUE, ROCKVILLE, MD, 20850 | | |
| NUMBER OF CLAIMS: | 49 | | |
| EXEMPLARY CLAIM: | 1 | | |
| NUMBER OF DRAWINGS: | 33 Drawing Page(s) | | |
| LINE COUNT: | 12881 | | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB Human TNF-gamma-alpha and TNF-gamma-beta polypeptides and DNA (RNA) encoding such polypeptides and a procedure for producing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing such polypeptides to inhibit cellular growth, for example in a tumor or cancer, for facilitating wound-healing, to provide resistance against infection, induce inflammatory activities, and stimulating the growth of certain cell types to treat diseases, for example restenosis. Also disclosed are diagnostic methods for detecting a mutation in the TNF-gamma-alpha and TNF-gamma-beta nucleic acid sequences or overexpression of the TNF-gamma-alpha and/or TNF-gamma-beta polypeptides. Antagonists against such polypeptides and their use as a therapeutic to treat cachexia, septic shock, cerebral malaria, inflammation, arthritis and graft-rejection are also disclosed.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 25 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2002:171934 USPATFULL
TITLE: TUMOR NECROSIS FACTOR-GAMMA
INVENTOR(S): YU, GUO-LIANG, SAN MATEO, CA, UNITED STATES
NI, JIAN, ROCKVILLE, MD, UNITED STATES
ROSEN, CRAIG A., LAYTONSVILLE, MD, UNITED STATES
ZHANG, JUN, BETHESDA, MD, UNITED STATES

| | NUMBER | KIND | DATE | |
|-----------------------|---|------|--------------|-----|
| PATENT INFORMATION: | US 2002090683 | A1 | 20020711 | <-- |
| | US 6824767 | B2 | 20041130 | |
| APPLICATION INFO.: | US 1999-246129 | A1 | 19990208 (9) | |
| RELATED APPLN. INFO.: | Continuation-in-part of Ser. No. US 1998-131237, filed on 7 Aug 1998, PENDING Continuation-in-part of Ser. No. US 1998-5020, filed on 9 Jan 1998, ABANDONED Continuation-in-part of Ser. No. US 1995-461246, filed on 5 Jun 1995, ABANDONED Continuation-in-part of Ser. No. WO 1994-US12880, filed on 7 Nov 1994, UNKNOWN | | | |

| NUMBER | DATE |
|--------|-------|
| ----- | ----- |

PRIORITY INFORMATION: US 1998-74047P 19980209 (60) <--
 DOCUMENT TYPE: Utility
 FILE SEGMENT: APPLICATION
 LEGAL REPRESENTATIVE: HUMAN GENOME SCIENCES INC, 9410 KEY WEST AVENUE,
 ROCKVILLE, MD, 20850
 NUMBER OF CLAIMS: 41
 EXEMPLARY CLAIM: 1
 NUMBER OF DRAWINGS: 30 Drawing Page(s)
 LINE COUNT: 6959

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB Human TNF-gamma-alpha and TNF-gamma-beta polypeptides and DNA (RNA) encoding such polypeptides and a procedure for producing such polypeptides by recombinant techniques is disclosed. Also disclosed are methods for utilizing such polypeptides to inhibit cellular growth, for example in a tumor or cancer, for facilitating wound-healing, to provide resistance against infection, induce inflammatory activities, and stimulating the growth of certain cell types to treat diseases, for example restenosis. Also disclosed are diagnostic methods for detecting a mutation in the TNF-gamma-alpha and TNF-gamma-beta nucleic acid sequences or overexpression of the TNF-gamma-alpha and TNF-gamma-beta polypeptides. Antagonists against such polypeptides and their use as a therapeutic to treat cachexia, septic shock, cerebral malaria, inflammation, arthritis and graft-rejection are also disclosed.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 26 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2002:346983 USPATFULL
 TITLE: Rin2, a novel inhibitor of Ras-mediated signaling
 INVENTOR(S): Tam, See-Ying, Mountain View, CA, United States
 Tsai, Mindy, Mountain View, CA, United States
 Galli, Stephen J., Portola Valley, CA, United States
 PATENT ASSIGNEE(S): Beth Israel Deaconess Medical Center, Palo Alto, CA,
 United States (U.S. corporation)
 The Board of Trustees of the Leland Stanford, Jr.,
 University, Palo Alto, CA, United States (U.S.
 corporation)

| | NUMBER | KIND | DATE |
|-----------------------|---|------|--------------|
| PATENT INFORMATION: | US 6500942 | B1 | 20021231 |
| APPLICATION INFO.: | US 2000-522955 | | 20000310 (9) |
| RELATED APPLN. INFO.: | Continuation of Ser. No. WO 1998-US19056, filed on 11 Sep 1998 Continuation-in-part of Ser. No. US 1997-942819, filed on 2 Oct 1997, now patented, Pat. No. US 5965707 | | |

| | NUMBER | DATE |
|-----------------------|--|-------------------|
| PRIORITY INFORMATION: | US 1997-58520P | 19970911 (60) <-- |
| DOCUMENT TYPE: | Utility | |
| FILE SEGMENT: | GRANTED | |
| PRIMARY EXAMINER: | Eyler, Yvonne | |
| ASSISTANT EXAMINER: | Nickol, Gary B. | |
| LEGAL REPRESENTATIVE: | Hamilton, Brook, Smith & Reynolds, P.C. | |
| NUMBER OF CLAIMS: | 45 | |
| EXEMPLARY CLAIM: | 1 | |
| NUMBER OF DRAWINGS: | 22 Drawing Figure(s); 18 Drawing Page(s) | |
| LINE COUNT: | 2376 | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB Novel gene, rin2, and encoded protein are disclosed which can inhibit the functional response induced by Ras-dependent signaling pathways are disclosed. Methods of inhibiting or enhancing Ras-dependent signaling and methods of treatment utilizing Rin2 are also disclosed.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 27 OF 34 USPATFULL on STN

ACCESSION NUMBER: 2002:181690 USPATFULL

TITLE: Substituted pyrazoles as p38 kinase inhibitors

INVENTOR(S): Anantanarayan, Ashok, Hainesville, IL, United States

Clare, Michael, Skokie, IL, United States

Collins, Paul W., Deerfield, IL, United States

Crich, Joyce Zuowu, Glenview, IL, United States

Devraj, Rajesh, Ballwin, MO, United States

Flynn, Daniel L., Clarkson Valley, MO, United States

Geng, Lifeng, Skokie, IL, United States

Graneto, Matthew J., Chesterfield, MO, United States

Hanau, Cathleen E., Chesterfield, MO, United States

Hanson, Gunnar J., Skokie, IL, United States

Hartmann, Susan J., Kirkwood, MO, United States

Hepperle, Michael, St. Charles, MO, United States

Huang, He, Chicago, IL, United States

Koszyk, Francis J., Prospect Heights, IL, United States

Liao, Shuyuan, Glen Ellyn, IL, United States

Metz, Suzanne, Chesterfield, MO, United States

Partis, Richard A., Evanston, IL, United States

Perry, Thao D., Chesterfield, MO, United States

Rao, Shashidhar N., St. Louis, MO, United States

Selness, Shaun Raj, Chesterfield, MO, United States

South, Michael S., St. Louis, MO, United States

Stealey, Michael A., Libertyville, IL, United States

Talley, John Jeffrey, St. Louis, MO, United States

Vazquez, Michael L., Ballwin, MO, United States

Weier, Richard M., Lake Bluff, IL, United States

Xi, Xiangdong, Evanston, IL, United States

Khanna, Ish K., Libertyville, IL, United States

Yu, Yi, Skokie, IL, United States

PATENT ASSIGNEE(S): G. D. Searle & Company, Skokie, IL, United States (U.S. corporation)

| | NUMBER | KIND | DATE |
|-----------------------|---|------|--------------|
| PATENT INFORMATION: | US 6423713 | B1 | 20020723 |
| APPLICATION INFO.: | US 2001-918481 | | 20010731 (9) |
| RELATED APPLN. INFO.: | Division of Ser. No. US 1998-196623, filed on 20 Nov 1998 Continuation-in-part of Ser. No. US 1998-83670, filed on 22 May 1998, now abandoned | | |

| | NUMBER | DATE |
|-----------------------|--|---------------|
| PRIORITY INFORMATION: | US 1997-47570P | 19970522 (60) |
| DOCUMENT TYPE: | Utility | |
| FILE SEGMENT: | GRANTED | |
| PRIMARY EXAMINER: | Solola, T. A. | |
| LEGAL REPRESENTATIVE: | Harness, Dickey & Pierce, P.L.C. | |
| NUMBER OF CLAIMS: | 88 | |
| EXEMPLARY CLAIM: | 1 | |
| NUMBER OF DRAWINGS: | 0 Drawing Figure(s); 0 Drawing Page(s) | |
| LINE COUNT: | 16941 | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB A class of pyrazole derivatives is described for use in treating p38 kinase mediated disorders. Compounds of particular interest are defined by Formula IA ##STR1##

wherein R.sup.1, R.sup.2, R.sup.3 and R.sup.4 are as described in the specification.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L24 ANSWER 28 OF 34 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2002:742981 HCAPLUS

DOCUMENT NUMBER: 138:53701

TITLE: p38 and Activating Transcription Factor-2 Involvement in Osteoblast Osmotic Response to Elevated Extracellular Glucose

AUTHOR(S): Zayzafoon, Majd; Botolin, Sergiu; McCabe, Laura R.

CORPORATE SOURCE: Department of Physiology, Michigan State University, East Lansing, MI, 48824, USA

SOURCE: Journal of Biological Chemistry (2002), 277(40), 37212-37218

CODEN: JBCHA3; ISSN: 0021-9258

PUBLISHER: American Society for Biochemistry and Molecular Biology

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Poorly controlled or untreated **type I diabetes**

mellitus is characterized by hyperglycemia and is associated with decreased bone mass and osteoporosis. We have demonstrated that osteoblasts are sensitive to hyperglycemia-associated osmotic stress and respond to elevated extracellular glucose or mannitol by increasing c-jun and collagen I expression. To determine whether MAPKs are involved in this response, MC3T3-E1 osteoblasts were treated with 16.5 mM glucose, mannitol, or contrast dye for 1 h. Immunoblotting of phosphorylated p38 demonstrated activation of p38 MAPK by hyperosmotic stress in vitro and in vivo. Activation peaked at 20 min, remained detectable after 24 h, and was protein kinase C-independent. Activating transcription factor-2 (ATF-2) activation followed the same pattern as phospho-p38. Transactivation of cAMP response element (CRE)- and c-jun promoter (containing a CRE-like element)-reporter constructs increased following hyperosmotic treatment. SB 203580 (a **p38 MAPK inhibitor**) blocked

ATF-2 phosphorylation, CRE transactivation, and c-jun promoter activation. Hyperosmotic activation of collagen I promoter activity was also inhibited by SB 203580, consistent with the involvement of c-jun in collagen I up-regulation. Therefore, we propose that hyperglycemia-induced increases in p38 MAPK activity and ATF-2 phosphorylation contribute to CRE activation and modulation of c-jun and collagen I expression in osteoblasts.

REFERENCE COUNT: 73 THERE ARE 73 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L24 ANSWER 29 OF 34 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2002:521726 HCAPLUS

DOCUMENT NUMBER: 137:214971

TITLE: Hyperketonemia increases tumor necrosis factor- α secretion in cultured U937 monocytes and type 1 diabetic patients and is apparently mediated by oxidative stress and cAMP deficiency

AUTHOR(S): Jain, Sushil K.; Kannan, Krishnaswamy; Lim, Gideon; McVie, Robert; Bocchini, Joseph A., Jr.

CORPORATE SOURCE: Department of Pediatrics, Louisiana State University
Health Sciences Center, Shreveport, LA, 71130, USA
SOURCE: Diabetes (2002), 51(7), 2287-2293
CODEN: DIAEAB; ISSN: 0012-1797
PUBLISHER: American Diabetes Association
DOCUMENT TYPE: Journal
LANGUAGE: English

AB An elevated blood level of tumor necrosis factor (TNF)- α is a validated marker of vascular inflammation, which can result in the development of vascular disease and atherosclerosis. This study examined the hypothesis that ketosis increases the TNF- α secretion, both in a cell culture model using U937 monocytes and in type 1 diabetic patients in vivo. U937 cells were cultured with ketone bodies (acetoacetate [AA] and β -hydroxybutyrate [BHB]) in the presence or absence of high levels of glucose in medium at 37° for 24 h. This study demonstrates the following points. First, hyperketonemic diabetic patients have significantly higher levels of TNF- α than normoketonemic diabetic patients ($P < 0.01$) and normal control subjects ($P < 0.01$). There was a significant correlation ($r = 0.36$, $P < 0.05$; $n = 34$) between ketosis and oxidative stress as well as between oxidative stress and TNF- α levels ($r = 0.47$, $P < 0.02$; $n = 34$) in the blood of diabetic patients. Second, ketone body AA treatment increases TNF- α secretion, increases oxygen radicals production, and lowers cAMP levels in U937 cells. However, BHB did not have any effect on TNF- α secretion or oxygen radicals production in U937 cells. Third, exogenous addition of dibutyryl

cAMP, endogenous stimulation of cAMP production by forskolin, and antioxidant N-acetylcysteine (NAC) prevented stimulation of TNF- α secretion caused by AA alone or with high glucose. Similarly, NAC prevented the elevation of TNF- α secretion and lowering of cAMP levels in H₂O₂-treated U937 cells. Fourth, the effect of AA on TNF- α secretion was inhibited by specific inhibitors of protein kinase A (H89), p38-mitogen-activated protein kinase (SB203580), and nuclear transcription factor (NF κ B) (NF κ B-SN50). This study demonstrates that hyperketonemia increases TNF- α secretion in cultured U937 monocytic cells and TNF- α levels in the blood of type 1 diabetic patients and is apparently mediated by AA-induced cellular oxidative stress and cAMP deficiency.

L24 ANSWER 30 OF 34 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2001:870187 HCAPLUS
DOCUMENT NUMBER: 136:132847
TITLE: A role for mitogen-activated protein kinases in the etiology of diabetic neuropathy
AUTHOR(S): Purves, Tertia; Middlemas, Alicia; Agthong, Sithiporn; Jude, Edward B.; Boulton, Andrew J. M.; Fernyhough, Paul; Tomlinson, David R.
CORPORATE SOURCE: Neuroscience Division, School of Biological Sciences, University of Manchester, Manchester, UK
SOURCE: FASEB Journal (2001), 15(13), 2508-2514
CODEN: FAJOEC; ISSN: 0892-6638
PUBLISHER: Federation of American Societies for Experimental Biology
DOCUMENT TYPE: Journal
LANGUAGE: English

AB The onset of diabetic neuropathy, a complication of diabetes mellitus, has been linked to poor glycemic control. We tested the hypothesis that the mitogen-activated protein kinases (MAPK) form transducers for the damaging effects of high glucose. In cultures of adult rat sensory neurons, high glucose activated JNK and p38 MAPK but did

not result in cell damage. However, oxidative stress activated ERK and p38 MAPKs and resulted in cellular damage. In the dorsal root ganglia of streptozotocin-induced diabetic rats (a model of **type I diabetes**), ERK and p38 were activated at 8 wk duration, followed by activation of JNK at 12 wk duration. We report activation of JNK and increases in total levels of p38 and JNK in sural nerve of type I and II diabetic patients. These data implicate MAPKs in the etiol. of diabetic neuropathy both via direct effects of glucose and via glucose-induced oxidative stress.

REFERENCE COUNT: 45 THERE ARE 45 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L24 ANSWER 31 OF 34 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2001:154506 HCAPLUS

DOCUMENT NUMBER: 135:370453

TITLE: Monocyte chemoattractant protein-1 is expressed in pancreatic islets from prediabetic NOD mice and in interleukin-1 β -exposed human and rat islet cells

AUTHOR(S): Chen, M.-C.; Proost, P.; Gysemans, C.; Mathieu, C.; Eizirik, D. L.

CORPORATE SOURCE: Gene Expression Unit, Diabetes Research Center, Vrije Universiteit Brussel, Brussels, Belg.

SOURCE: Diabetologia (2001), 44(3), 325-332

CODEN: DBTGAI; ISSN: 0012-186X

PUBLISHER: Springer-Verlag

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Monocyte chemoattractant protein-1 (MCP-1) attracts monocytes and T lymphocytes, and could thus contribute to mononuclear cell infiltration in type I (insulin-dependent) **diabetes** mellitus. Cytokines induce MCP-1 mRNA expression in pancreatic rat β cells. To investigate this issue, the authors analyzed the signal transduction for IL-1 β -induced MCP-1 expression in rat β cells and in vitro MCP-1 mRNA expression and protein release by human islets as well as in vivo islet MCP-1 mRNA expression in prediabetic non-obese diabetic mice. Fluorescence-activated cell sorting-purified rat β cells were cultured for 6 h with IL-1 β (30 U/mL) or MAPK inhibitors or both. Human islets were cultured for 6-72 h with the cytokines IL-1 β , IFN- γ , or the inducible nitric oxide synthase (iNOS) inhibitor NG-methyl-L-arginine or both. The authors measured MCP-1 mRNA by RT-PCR and protein by ELISA. The MCP-1 mRNA expression in islets from male and female non-obese diabetic mice (2-12 wk of age) was measured by real time reverse transcription-polymerase chain reaction (RT-PCR). Interleukin-1 β induced MCP-1 mRNA expression in rat β cells, with a maximum induction after 6 h. A combination of p38 and ERK1/2 inhibitors decreased MCP-1 expression by 70%. IL-1 β induced both MCP-1 mRNA expression and a 3-fold increase in medium MCP-1 protein accumulation in human islet cells. This effect was not prevented by iNOS blockers. In vivo there was an age-related increase in MCP-1 mRNA expression in islets from male and female non-obese diabetic mice, reaching a peak at 8 wk. Thus, in rat and human islet cells MCP-1 mRNA is induced by IL-1 β . Both ERK1/2 and p38 MAPK, but not nitric oxide, contribute to MCP-1 expression. In non-obese diabetic mice MCP-1 mRNA expression increases with age, peaking at the early phases of insulinitis. The production of MCP-1 by pancreatic beta cells could contribute to the recruitment of mononuclear cells into pancreatic islets in early **Type I diabetes**.

REFERENCE COUNT: 56 THERE ARE 56 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L24 ANSWER 32 OF 34 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2001:506695 HCAPLUS
DOCUMENT NUMBER: 136:83924
TITLE: Increased cytokine-induced cytotoxicity of pancreatic islet cells from transgenic mice expressing the Src-like tyrosine kinase GTK
AUTHOR(S): Anneren, Cecilia; Welsh, Michael
CORPORATE SOURCE: Department of Medical Cell Biology, Uppsala University, Uppsala, Swed.
SOURCE: Molecular Medicine (Baltimore, MD, United States) (2001), 7(5), 301-310
CODEN: MOMEF3; ISSN: 1076-1551
PUBLISHER: Johns Hopkins University Press
DOCUMENT TYPE: Journal
LANGUAGE: English

AB The loss of β cells in **type 1 diabetes** may involve protein kinases because they control cell growth, differentiation, and survival. Previous studies have revealed that GTK, a Src-like protein tyrosine kinase expressed in β cells (also named Bsk/Iyk), regulates multiple responses including growth and survival of rat insulinoma cells (RINm5F) and differentiation of neuronal PC12 cells. In the present study, the authors have generated a transgenic mouse expressing a kinase active GTK mutant (GTK-Y504F) under the control of the rat insulin I promoter to establish a role of GTK in β cells. Control and GTK-transgenic CBA mice were used for determination of *in vivo* glucose tolerance and the relative insulin-pos. area. Isolated islets from both groups were cultured in the absence and presence of cytokines and insulin secretion, viability and protein expression were assessed. The β -cell mass of GTK-transgenic mice was increased as a consequence of a larger pancreas and an increased relative β -cell area. Islets isolated from the transgenic animals exhibited an enhanced glucose-induced insulin release and reduced viability in response to cytokines that could not be explained by higher levels of NO compared with control islets. Extra-cellular signal-regulated kinase (ERK) 1/2, p38 mitogen-activated protein kinase (MAPK), c-Jun NH2-terminal kinase (JNK), and Akt were all activated by cytokines, but GTK-transgenic islets contained higher basal levels of phosphorylated ERK1/2 and lower basal levels of phosphorylated p38 compared with the control islets. The total amount of activated MAPKs was, however, higher in the cytokine-stimulated transgenic islets compared with the control islets due to increased levels of phospho-ERK1/2. Moreover, the proline-rich tyrosine kinase (PYK) 2 (also named RAFTK/CAK β /CADTK) levels were elevated in response to a 24-h exposure to cytokines in control islets but not in the GTK-transgenic islets. These results suggest that although GTK increases the β -cell mass, it also enhances islet cell death in response to cytokines and may thus be involved in the β -cell damage in **type 1 diabetes**.

REFERENCE COUNT: 45 THERE ARE 45 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L24 ANSWER 33 OF 34 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2000:627254 HCAPLUS
DOCUMENT NUMBER: 133:295222
TITLE: Activation of extracellular signal-regulated kinase (ERK) 1/2 contributes to cytokine-induced apoptosis in purified rat pancreatic β -cells
AUTHOR(S): Pavlovic, Dejan; Andersen, Nina Aa.; Mandrup-Poulsen, Thomas; Eizirik, Decio L.
CORPORATE SOURCE: Gene Expression Unit Diabetes Research Center, Vrije Universiteit Brussel, Brussels, B-1090, Belg.

SOURCE: European Cytokine Network (2000), 11(2),
267-274
CODEN: ECYNEJ; ISSN: 1148-5493
PUBLISHER: John Libbey Eurotext
DOCUMENT TYPE: Journal
LANGUAGE: English

AB Cytokines may contribute to β -cell apoptosis in the early stages of **type 1 diabetes** mellitus. It has been reported recently that interleukin- 1β (IL- 1β) induces activation of the mitogen-activated protein kinases (MAPK) p38 and ERK1/2 in neonatal rat islets. Since these kinases may participate in cytokine-induced apoptosis, we evaluated whether cytokines induce activation of MAPKs in FACS-purified primary rat β -cells, and whether blockers of p38 and/or ERK1/2 prevent β -cell death. IL- 1β , but not interferon- γ (IFN- γ), caused phosphorylation of the substrates Elk-1, ATF-2 and hsp25, and the phosphorylation of both Elk-1 and hsp25 were decreased by the p38 blocker SB203580 (p38i) and the MAPK/ERK blocker PD 098059 (MEKi). When added together, p38i and MEKi decreased IL- 1β -induced nitrite production over 24 h by 60%, but did not affect IL- 1β -induced manganese superoxide dismutase (MnSOD) mRNA expression. To test the effects of MAPK inhibitors on β -cell death by necrosis or apoptosis, these cells were exposed for 6 or 9 days to IL- 1β + IFN- γ . This treatment induced cell death, mostly by apoptosis. The MEKi, but not the p38i, significantly decreased cytokine-induced apoptosis, thus decreasing the total number of dead cells. This protection was only partial, suggesting that ERK1/2 activation is not the only mechanism by which cytokines induce β -cell apoptosis. We conclude that IL- 1β induces activation of both p38 and ERK1/2, and that ERK1/2 contributes to the pro-apoptotic effects of the cytokine in primary β -cells.

REFERENCE COUNT: 42 THERE ARE 42 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L24 ANSWER 34 OF 34 USPATFULL on STN

ACCESSION NUMBER: 1999:125043 USPATFULL
TITLE: Rin2, a novel inhibitor of Ras-mediated signaling
INVENTOR(S): Tam, See-Ying, Wellesley, MA, United States
Tsai, Mindy, Wellesley, MA, United States
Galli, Stephen J., Winchester, MA, United States
PATENT ASSIGNEE(S): Beth Israel Deaconess Medical Center, Boston, MA,
United States (U.S. corporation)

| | NUMBER | KIND | DATE | |
|---------------------|----------------|------|--------------|-----|
| PATENT INFORMATION: | US 5965707 | | 19991012 | <-- |
| APPLICATION INFO.: | US 1997-942819 | | 19971002 (8) | |

| | NUMBER | DATE | |
|-----------------------|---|---------------|-----|
| PRIORITY INFORMATION: | US 1997-58520P | 19970911 (60) | <-- |
| DOCUMENT TYPE: | Utility | | |
| FILE SEGMENT: | Granted | | |
| PRIMARY EXAMINER: | Hutzell, Paula K. | | |
| ASSISTANT EXAMINER: | Sun-Hoffman, Lin | | |
| LEGAL REPRESENTATIVE: | Hamilton, Brook, Smith & Reynolds, P.C. | | |
| NUMBER OF CLAIMS: | 8 | | |
| EXEMPLARY CLAIM: | 1 | | |
| NUMBER OF DRAWINGS: | 10 Drawing Figure(s); 8 Drawing Page(s) | | |
| LINE COUNT: | 1882 | | |

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB Novel gene, rin2, and encoded protein are disclosed which can inhibit

the functional response induced by Ras-dependent signaling pathways are disclosed. Methods of inhibiting or enhancing Ras-dependent signaling and methods of treatment utilizing Rin2 are also disclosed.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

TEXT SEARCH - MEDLINE BIOSIS EMBASE JAPIO JICST

=> d que stat l22

L17 2 SEA FILE=REGISTRY ABB=ON ("P38 MAPK"/CN OR "P38 MAPK (BIOMPHAL
ARIA GLABRATA STRAIN BS90 HEMOCYTE)"/CN OR "P38 MAPK INHIBITOR"
/CN)

L18 10002 SEA FILE=HCAPLUS ABB=ON L17 OR P38(W)?MAPK?(W)?INHIBIT?

L19 320 SEA FILE=HCAPLUS ABB=ON L18 AND ?DIABETES?

L20 25 SEA FILE=HCAPLUS ABB=ON L19 AND (TYPE(W)(1 OR I)(W)?DIABETES?)

L22 23 SEA L20

=> d ibib abs l22 1-23

L22 ANSWER 1 OF 23 MEDLINE on STN

ACCESSION NUMBER: 2002493287 MEDLINE

DOCUMENT NUMBER: PubMed ID: 12149242

TITLE: P38 and activating transcription factor-2 involvement in
osteoblast osmotic response to elevated extracellular
glucose.

AUTHOR: Zayzafoon Majd; Botolin Sergiu; McCabe Laura R

CORPORATE SOURCE: Department of Physiology, Michigan State University, East
Lansing, Michigan 48824, USA.

SOURCE: The Journal of biological chemistry, (2002 Oct 4) Vol. 277,
No. 40, pp. 37212-8. Electronic Publication: 2002-07-30.
Journal code: 2985121R. ISSN: 0021-9258.

PUB. COUNTRY: United States

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English

FILE SEGMENT: Priority Journals

ENTRY MONTH: 200211

ENTRY DATE: Entered STN: 1 Oct 2002
Last Updated on STN: 5 Jan 2003
Entered Medline: 20 Nov 2002

AB Poorly controlled or untreated **type I diabetes**
mellitus is characterized by hyperglycemia and is associated with
decreased bone mass and osteoporosis. We have demonstrated that
osteoblasts are sensitive to hyperglycemia-associated osmotic stress and
respond to elevated extracellular glucose or mannitol by increasing c-jun
and collagen I expression. To determine whether MAPKs are involved in
this response, MC3T3-E1 osteoblasts were treated with 16.5 mM glucose,
mannitol, or contrast dye for 1 h. Immunoblotting of phosphorylated p38
demonstrated activation of p38 MAPK by hyperosmotic stress in vitro and in
vivo. Activation peaked at 20 min, remained detectable after 24 h, and
was protein kinase C-independent. Activating transcription factor-2
(ATF-2) activation followed the same pattern as phospho-p38.
Transactivation of cAMP response element (CRE)- and c-jun promoter
(containing a CRE-like element)-reporter constructs increased following
hyperosmotic treatment. SB 203580 (a **p38 MAPK**
inhibitor) blocked ATF-2 phosphorylation, CRE transactivation, and
c-jun promoter activation. Hyperosmotic activation of collagen I promoter
activity was also inhibited by SB 203580, consistent with the involvement
of c-jun in collagen I up-regulation. Therefore, we propose that
hyperglycemia-induced increases in p38 MAPK activity and ATF-2
phosphorylation contribute to CRE activation and modulation of c-jun and
collagen I expression in osteoblasts.

L22 ANSWER 2 OF 23 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN

ACCESSION NUMBER: 2006:453398 BIOSIS

DOCUMENT NUMBER: PREV200600449427

TITLE: Insulin signal mimicry as a mechanism for the insulin-like

effects of vanadium.
AUTHOR(S): Mehdi, Mohamad Z.; Pandey, Sanjay K.; Theberge,
Jean-Francois; Srivastava, Ashok K. [Reprint Author]
CORPORATE SOURCE: CHU Montreal, Hotel Dieu, Lab Cell Signaling, Res Ctr,
Montreal, PQ, Canada
ashok.srivastava@umontreal.ca
SOURCE: Cell Biochemistry and Biophysics, (2006) Vol. 44, No. 1,
pp. 73-81.
ISSN: 1085-9195.
DOCUMENT TYPE: Article
General Review; (Literature Review)
LANGUAGE: English
ENTRY DATE: Entered STN: 13 Sep 2006
Last Updated on STN: 13 Sep 2006

AB Among several metals, vanadium has emerged as an extremely potent agent with insulin-like properties. These insulin-like properties have been demonstrated in isolated cells, tissues, different animal models of type I and type II **diabetes** as well as a limited number of human subjects. Vanadium treatment has been found to improve abnormalities of carbohydrate and lipid metabolism and of gene expression in rodent models of **diabetes**. In isolated cells, it enhances glucose transport, glycogen and lipid synthesis, and inhibits gluconeogenesis and lipolysis. The molecular mechanism responsible for the insulin-like effects of vanadium compounds have been shown to involve the activation of several key components of insulin-signaling pathways that include the mitogen-activated-protein kinases (MAPKs) extracellular signal-regulated kinase 1/2 (ERK1/2) and p38NIPAK, and phosphatidylinositol 3-kinase (PI3-K)/protein kinase B (PKB). It is interesting that the vanadium effect on these signaling systems is independent of insulin receptor protein tyrosine kinase activity, but it is associated with enhanced tyrosine phosphorylation of insulin receptor substrate-1. These actions seem to be secondary to vanadium-induced inhibition of protein tyrosine phosphatases. Because MAPK and PI3-K/PKB pathways are implicated in mediating the mitogenic and metabolic effects of insulin, respectively, it is plausible that mimicry of these pathways by vanadium serves as a mechanism for its insulin-like responses.

L22 ANSWER 3 OF 23 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN
ACCESSION NUMBER: 2006:412874 BIOSIS
DOCUMENT NUMBER: PREV200600415391
TITLE: Elevated glucose and **diabetes** promote
interleukin-12 cytokine gene expression in mouse
macrophages.
AUTHOR(S): Wen, Yeshao; Gu, Jiali; Li, Shu-Lian; Reddy, Marpadga A.;
Natarajan, Rama; Nadler, Jerry L. [Reprint Author]
CORPORATE SOURCE: Univ Virginia, Diabet and Hormone Ctr, Charlottesville, VA
22908 USA
jln2n@virginia.edu
SOURCE: Endocrinology, (MAY 2006) Vol. 147, No. 5, pp. 2518-2525.
CODEN: ENDOAO. ISSN: 0013-7227.
DOCUMENT TYPE: Article
LANGUAGE: English
ENTRY DATE: Entered STN: 23 Aug 2006
Last Updated on STN: 23 Aug 2006

AB Inflammation is emerging as an important mechanism for microand macrovascular complication of **diabetes**. The macrophage plays a key role in the chronic inflammatory response in part by generating particular cytokines. IL-1 beta, IL-6, IL12, IL-18, TNF alpha, and interferon-gamma are produced primarily in macrophages and have been associated with accelerated atherosclerosis and altered vascular wall

function. In this study, we evaluated the effect and mechanism of high glucose (HG) on gene expression of these cytokines in mouse peritoneal macrophages (MPM). HG led to a 2-fold increase in the mRNA expression of these cytokines, with IL-12 showing the highest activation (5.4-fold) in a time-dependent (3-12 h) and dosedependent (10, 17.5, and 25 mmol/liter) manner. The effects were specific to HG because mannitol and 3-O-methyl-glucose had no effect on cytokine mRNA expression. HG also increased IL-12 protein accumulation from MPM. We also explored the role of induced and spontaneous diabetes on inflammatory cytokine expression in MPM. Increases in expression in MPM of multiple inflammatory cytokines, including a 20-fold increase in IL-12 mRNA, were observed in streptozotocin- induced type 1 diabetic mice as well as type 2 diabetic db/db mice, suggesting that cytokine gene expression is increased by hyperglycemia in vivo. We next explored potential mechanisms of HG-induced increases in IL-12 mRNA. HGincreased the activity of protein kinase C, p38MAPK (p38), c-Jun terminal kinase, and inhibitory-kappa B kinase in MPM. Furthermore, inhibitors of these signaling pathways significantly reduced HG-induced IL-12 mRNA expression in MPM. These results provide evidence for a potentially important mechanism linking elevated glucose and diabetes to inflammation.

L22 ANSWER 4 OF 23 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN
 ACCESSION NUMBER: 2006:351802 BIOSIS
 DOCUMENT NUMBER: PREV200600364780
 TITLE: Defective NF-kappaB activation in response to LPS in
 type 1 diabetes dendritic
 cells.
 AUTHOR(S): Mollah, Z. U. A. [Reprint Author]; Moore, C.; Sullivan, B.;
 Peng, J.; Phillips, K.; Cardinal, J.; Prins, J.; Thomas, R.
 CORPORATE SOURCE: Ctr Immunol and Canc Res, Buranda, Qld, Australia
 SOURCE: Tissue Antigens, (NOV 2005) Vol. 66, No. 5, pp. 496.
 Meeting Info.: 35th Annual Scientific Meeting of the
 Australasian-Society-for-Immunology/14th International HLA
 and Immunogenetics Workshops. Melbourne, AUSTRALIA.
 November 29 -December 02, 2005. Australasian Soc Immunol.
 CODEN: TSANA2. ISSN: 0001-2815.
 DOCUMENT TYPE: Conference; (Meeting)
 Conference; Abstract; (Meeting Abstract)
 LANGUAGE: English
 ENTRY DATE: Entered STN: 19 Jul 2006
 Last Updated on STN: 19 Jul 2006

L22 ANSWER 5 OF 23 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN
 ACCESSION NUMBER: 2006:179259 BIOSIS
 DOCUMENT NUMBER: PREV200600161409
 TITLE: Glucose-induced reactive oxygen species cause apoptosis of
 podocytes and podocyte depletion at the onset of diabetic
 nephropathy.
 AUTHOR(S): Susztak, Katalin; Raff, Amanda C.; Schiffer, Mario;
 Boettinger, Erwin P. [Reprint Author]
 CORPORATE SOURCE: Mt Sinai Sch Med, Div Nephrol, Dept Med, 1 Gustave L Levy
 Pl, New York, NY 10029 USA
 ksusztak@aecom.yu.edu
 SOURCE: Diabetes, (JAN 2006) Vol. 55, No. 1, pp. 225-233.
 CODEN: DIAEAZ. ISSN: 0012-1797.
 DOCUMENT TYPE: Article
 LANGUAGE: English
 ENTRY DATE: Entered STN: 9 Mar 2006
 Last Updated on STN: 9 Mar 2006
 AB Diabetic nephropathy is the most common cause of end-stage renal disease

in the U.S. Recent studies demonstrate that loss of podocytes is an early feature of diabetic nephropathy that predicts its progressive course. Cause and consequences of podocyte loss during early diabetic nephropathy remain poorly understood. Here, we demonstrate that podocyte apoptosis increased sharply with onset of hyperglycemia in Ins2(Akita) (Akita) mice with **type 1 diabetes** and Lepr(db/db) (db/db) mice with obesity and type 2 **diabetes**. Podocyte apoptosis coincided with the onset of urinary albumin excretion (UAE) and preceded significant losses of podocytes in Akita (37% reduction) and db/db (27% reduction) mice. Increased extracellular glucose (30 mmol/l) rapidly stimulated generation of intracellular reactive oxygen species (ROS) through NADPH oxidase and mitochondrial pathways and led to activation of proapoptotic p38 mitogen-activated protein kinase and caspase 3 and to apoptosis of conditionally immortalized podocytes in vitro. Chronic inhibition of NADPH oxidase prevented podocyte apoptosis and ameliorated podocyte depletion, UAE, and mesangial matrix expansion in db/db mice. In conclusion, our results demonstrate for the first time that glucose-induced ROS production initiates podocyte apoptosis and podocyte depletion in vitro and in vivo and suggest that podocyte apoptosis/depletion represents a novel early pathomechanism(s) leading to diabetic nephropathy in murine type 1 and type 2 diabetic models.

L22 ANSWER 6 OF 23 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN
ACCESSION NUMBER: 2006:155490 BIOSIS
DOCUMENT NUMBER: PREV200600163314
TITLE: The inflammatory chemokine, interferon-gamma-inducible protein (IP-10), is induced in monocytes by ligands of the receptor for advanced glycation end products (RAGE).
AUTHOR(S): Shanmugam, Narkunaraja [Reprint Author]; Ransohoff, Richard M.; Natarajan, Rama
SOURCE: Diabetes, (JUN 2004) Vol. 53, No. Suppl. 2, pp. A450. Meeting Info.: 64th Annual Meeting of the American-Diabetes-Association. Orlando, FL, USA. June 04 -08, 2004. Amer Diabet Assoc. CODEN: DIAEAZ. ISSN: 0012-1797.
DOCUMENT TYPE: Conference; (Meeting)
Conference; Abstract; (Meeting Abstract)
LANGUAGE: English
ENTRY DATE: Entered STN: 9 Mar 2006
Last Updated on STN: 9 Mar 2006

L22 ANSWER 7 OF 23 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN
ACCESSION NUMBER: 2005:474184 BIOSIS
DOCUMENT NUMBER: PREV200510274945
TITLE: Effects of sodium tungstate in Min6 beta-cells: potential implications for its antidiabetic action.
AUTHOR(S): Piquer, S. [Reprint Author]; Barcelo-Batllori, S.; Julia, M.; Gomis, R.
CORPORATE SOURCE: Hosp Clin Barcelona, Lab Expt Diabet, Barcelona, Spain
SOURCE: Diabetologia, (AUG 2004) Vol. 47, No. Suppl. 1, pp. A169. Meeting Info.: 40th Annual Meeting of the European-Association-for-the-Study-of-Diabetes. Munich, GERMANY. September 05 -09, 2004. European Assoc Study Diabetes. CODEN: DBTGJ. ISSN: 0012-186X.
DOCUMENT TYPE: Conference; (Meeting)
Conference; Abstract; (Meeting Abstract)
LANGUAGE: English
ENTRY DATE: Entered STN: 16 Nov 2005
Last Updated on STN: 16 Nov 2005

L22 ANSWER 8 OF 23 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN
 ACCESSION NUMBER: 2005:402453 BIOSIS
 DOCUMENT NUMBER: PREV200510189321
 TITLE: Selective inhibition of Fc epsilon RI-mediated mast cell
 activation by a truncated variant of Cbl-b related to the
 rat model of **type 1 diabetes**
 mellitus.
 AUTHOR(S): Qu, Xiujuan; Miah, S. M. Shahjahan; Hatani, Tomoko;
 Okazaki, Mami; Hori-Tamura, Naoko; Yamamura, Hirohei;
 Hotta, Hak; Sada, Kiyonao [Reprint Author]
 CORPORATE SOURCE: Kobe Univ, Grad Sch Med, Dept Genome Sci, Div Proteom,
 Kobe, Hyogo 6500017, Japan
 ksada@med.kobe-u.ac.jp
 SOURCE: Journal of Biochemistry (Tokyo), (JUN 2005) Vol. 137, No.
 6, pp. 711-720.
 CODEN: JOBIAO. ISSN: 0021-924X.
 DOCUMENT TYPE: Article
 LANGUAGE: English
 ENTRY DATE: Entered STN: 5 Oct 2005
 Last Updated on STN: 5 Oct 2005

AB Ubiquitin-protein ligase Cbl-b negatively regulates high affinity IgE
 receptor (Fc epsilon RI)-mediated degranulation and cytokine gene
 transcription in mast cells. In this study, we have examined the role of
 a truncated variant of Cbl-b related to the rat model of **type**
1 diabetes mellitus using the mast cell signaling model.
 Overexpression of the truncated Cbl-b that lacks the C-terminal region did
 not suppress the activation of proximal and distal signaling molecules
 leading to degranulation. Fc epsilon CRI-mediated tyrosine
 phosphorylation of Syk, Gab2, and phospholipase C-gamma 1, and activation
 of c-Jun N-terminal kinase (JNK), extracellular signal-regulated kinase
 (ERK), p38 mitogen-activated protein kinase (MAP kinase), and inhibitor of
 nuclear factor kappa B kinase (IKK), and generation of Rac1 are unaffected
 in cells overexpressing the truncated Cbl-b in the lipid raft. On the
 other hand, Fc epsilon RI-mediated transcriptional activation of nuclear
 factor of activated T cells (NFAT), and transcription of interleukin-3
 (IL-3) and IL-4 mRNA are inhibited by overexpression of the truncated
 variant of Cbl-b. This suppression parallels the re-compartmentalization
 of specific effector molecules in the lipid raft. These structural and
 functional analyses reveal the mechanism underlying the selective
 inhibition of cellular signaling by the truncated variant of Cbl-b related
 to insulin-dependent **diabetes** mellitus.

L22 ANSWER 9 OF 23 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN
 ACCESSION NUMBER: 2005:123865 BIOSIS
 DOCUMENT NUMBER: PREV200500129411
 TITLE: C-peptide stimulates ERK1/2 and JNK MAP kinases via
 activation of protein kinase C in human renal tubular
 cells.
 AUTHOR(S): Zhong, Z.; Davidescu, A.; Ehren, I.; Ekberg, K.; Jornvall,
 H.; Wahren, J.; Chibalin, A. V. [Reprint Author]
 CORPORATE SOURCE: Dept Surg SciSect Integrat Physiol, Karolinska Inst, von
 Eulers Vag 4,4 Tr, S-17177, Stockholm, Sweden
 Alexander.Chibalin@kirurgi.ki.se
 SOURCE: Diabetologia, (January 2005) Vol. 48, No. 1, pp. 187-197.
 print.
 CODEN: DBTGAI. ISSN: 0012-186X.
 DOCUMENT TYPE: Article
 LANGUAGE: English
 ENTRY DATE: Entered STN: 1 Apr 2005

Last Updated on STN: 1 Apr 2005

AB Aims/hypothesis: Accumulating evidence indicates that replacement of C-peptide in **type 1 diabetes** ameliorates nerve and kidney dysfunction, but the molecular mechanisms involved are incompletely understood. C-peptide shows specific binding to a G-protein-coupled membrane binding site, resulting in Ca^{2+} influx, activation of mitogen-activated protein kinase signalling pathways, and stimulation of Na^{+} , K^{+} -ATPase and endothelial nitric oxide synthase. This study examines the intracellular signalling pathways activated by C-peptide in human renal tubular cells. Methods: Human renal tubular cells were cultured from the outer cortex of renal tissue obtained from patients undergoing elective nephrectomy. Extracellular-signal-regulated kinase 1/2 (ERK1/2), c-Jun N-terminal kinase (JNK) and Akt/protein kinase B (PKB) activation was determined using phospho-specific antibodies. Protein kinase C (PKC) and RhoA activation was determined by measuring their translocation to the cell membrane fraction using isoform-specific antibodies. Results: Human C-peptide increases phosphorylation of ERK1/2 and Akt/PKB in a concentration- and time-dependent manner in renal tubular cells. The C-terminal pentapeptide of C-peptide is equipotent with the full-length C-peptide, whereas scrambled C-peptide has no effect. C-peptide stimulation also results in phosphorylation of JNK, but not of p38 mitogen-activated protein kinase. MEK1/2 inhibitor PD98059 blocks the C-peptide effect on ERK1/2 phosphorylation. C-peptide causes specific translocation of PKC isoforms delta and epsilon to the membrane fraction in tubular cells. All stimulatory effects of C-peptide were abolished by pertussis toxin. The isoform-specific PKC-delta inhibitor rottlerin and the broad-spectrum PKC inhibitor GF109203X both abolish the C-peptide effect on ERK1/2 phosphorylation. C-peptide stimulation also causes translocation of the small GTPase RhoA from the cytosol to the cell membrane. Inhibition of phospholipase C abolished the stimulatory effect of C-peptide on phosphorylation of ERK1/2, JNK and PKC-delta. Conclusions/interpretation: C-peptide signal transduction in human renal tubular cells involves the activation of phospholipase C and PKC-delta and PKC-epsilon, as well as RhoA, followed by phosphorylation of ERK1/2 and JNK, and a parallel activation of Akt.

L22 ANSWER 10 OF 23 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN

ACCESSION NUMBER: 2004:425620 BIOSIS
DOCUMENT NUMBER: PREV200400422623
TITLE: Abnormal p38 mitogen-activated protein kinase signalling in human and experimental diabetic nephropathy.
AUTHOR(S): Adhikary, L.; Chow, F.; Nikolic-Paterson, D. J.; Stambe, C.; Dowling, J.; Atkins, R. C.; Tesch, G. H. [Reprint Author]
CORPORATE SOURCE: Dept Nephrol, Monash Med Ctr, 246 Clayton Rd, Clayton, Vic, 3168, Australia
gtesch@hotmail.com
SOURCE: Diabetologia, (July 2004) Vol. 47, No. 7, pp. 1210-1222. print.
CODEN: DBTG AJ. ISSN: 0012-186X.
DOCUMENT TYPE: Article
LANGUAGE: English
ENTRY DATE: Entered STN: 3 Nov 2004
Last Updated on STN: 3 Nov 2004

AB Aims/hypothesis. Inflammation and fibrosis are pathological mechanisms that are partially regulated by cell signalling through the p38 mitogen-activated protein kinase (MAPK) pathway. Elements of the diabetic milieu such as high glucose and advanced glycation end-products induce activation of this pathway in renal cells. Therefore, we examined whether

p38 MAPK signalling is associated with the development of human and experimental diabetic nephropathy. Methods. Immunostaining identified phosphorylated (active) p38 MAPK in human biopsies with no abnormality (n=6) and with Type 2 diabetic nephropathy (n=12). Changes in kidney levels of phosphorylated p38 were assessed by immunostaining and western blotting in mice with streptozotocin-induced **Type 1 diabetes** that had been killed after 0.5, 2, 3, 4 and 8 months, and in Type 2 diabetic db/db mice at 2, 4, 6 and 8 months of age. Results. Phosphorylated p38 was detected in some intrinsic cells in normal human kidney, including podocytes, cortical tubules and occasional interstitial cells. Greater numbers of these phosphorylated p38+ cells were observed in diabetic patients, and phosphorylated p38 was identified in accumulating interstitial macrophages and myofibroblasts. A similar pattern of p38 activation was observed in both mouse models of **diabetes**. In mice, kidney levels of phosphorylated p38 increased (2-6 fold) following the onset of Type 1 and Type 2 **diabetes**. In both mouse models, interstitial phosphorylated p38+ cells were associated with hyperglycaemia, increased HbA1c levels and albuminuria. Further assessment of streptozotocin-induced diabetic nephropathy showed that interstitial phosphorylated p38+ cells correlated with interstitial fibrosis (myofibroblasts, collagen). Conclusions/interpretation. Increased p38 MAPK signalling is a feature of human and experimental diabetic nephropathy. Time course studies in mouse models suggest that phosphorylation of p38 plays a pathological role, particularly in the development of interstitial fibrosis.

L22 ANSWER 11 OF 23 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN

ACCESSION NUMBER: 2004:178316 BIOSIS
 DOCUMENT NUMBER: PREV200400179575
 TITLE: Molecular mechanisms of high glucose-induced cyclooxygenase-2 expression in monocytes.
 AUTHOR(S): Shanmugam, Narkunaraja; Gaw Gonzalo, Irene T.; Natarajan, Rama [Reprint Author]
 CORPORATE SOURCE: Department of Diabetes, Beckman Research Institute of City of Hope, 1500 East Duarte Rd., Duarte, CA, 91010, USA
 rnatarajan@coh.org
 SOURCE: Diabetes, (March 2004) Vol. 53, No. 3, pp. 795-802. print. ISSN: 0012-1797 (ISSN print).
 DOCUMENT TYPE: Article
 LANGUAGE: English
 ENTRY DATE: Entered STN: 31 Mar 2004
 Last Updated on STN: 31 Mar 2004

AB The cyclooxygenase (COX)-2 enzyme has been implicated in the pathogenesis of several inflammatory diseases. However, its role in diabetic vascular disease is unclear. In this study, we evaluated the hypothesis that diabetic conditions can induce COX-2 in monocytes. High glucose treatment of THP-1 monocytic cells led to a significant three- to fivefold induction of COX-2 mRNA and protein expression but not COX-1 mRNA. High glucose-induced COX-2 mRNA was blocked by inhibitors of nuclear factor-kappaB (NF-kappaB), protein kinase C, and p38 mitogen-activated protein kinase. In addition, an antioxidant and inhibitors of mitochondrial superoxide, NADPH oxidase, and glucose metabolism to glucosamine also blocked high glucose-induced COX-2 expression to varying degrees. High glucose significantly increased transcription from a human COX-2 promoter-luciferase construct (twofold, $P < 0.001$). Promoter deletion analyses and inhibition of transcription by NF-kappaB superrepressor and cAMP-responsive element binding (CREB) mutants confirmed the involvement of NF-kappaB and CREB transcription factors in high glucose-induced COX-2 regulation. In addition, isolated peripheral

blood monocytes from type 1 and type 2 diabetic patients had high levels of COX-2 mRNA, whereas those from normal volunteers showed no expression. These results show that high glucose and **diabetes** can augment inflammatory responses by upregulating COX-2 via multiple signaling pathways, leading to monocyte activation relevant to the pathogenesis of **diabetes** complications.

L22 ANSWER 12 OF 23 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN

ACCESSION NUMBER: 2004:130132 BIOSIS

DOCUMENT NUMBER: PREV200400115748

TITLE: The specific p38 mitogen-activated protein kinase pathway inhibitor FR167653 keeps insulinitis benign in nonobese diabetic mice.

AUTHOR(S): Ando, Hitoshi; Kurita, Seiichiro; Takamura, Toshinari [Reprint Author]

CORPORATE SOURCE: Department of Endocrinology and Metabolism, Kanazawa University Graduate School of Medical Science, 13-1 Takara-machi, Kanazawa, Ishikawa, 920-8641, Japan
tt@medf.m.kanazawa-u.ac.jp

SOURCE: Life Sciences, (February 20 2004) Vol. 74, No. 14, pp. 1817-1827. print.

ISSN: 0024-3205 (ISSN print).

DOCUMENT TYPE: Article

LANGUAGE: English

ENTRY DATE: Entered STN: 3 Mar 2004

Last Updated on STN: 3 Mar 2004

AB The p38 mitogen-activated protein kinase (MAPK) pathway is important in Th1 immunity, macrophage activation, and apoptosis. Since they may be associated with beta-cell destruction during the development of **type 1 diabetes**, we investigated the role of the p38 MAPK pathway in female nonobese diabetic (NOD) mice. Phosphorylated p38 MAPK was observed immunohistochemically in CD4+ cells that had infiltrated into the islets and part of beta-cells, increasing in proportion to the severity of insulinitis. Continuous oral administration of 0.08% FR167653, a specific p38 MAPK pathway inhibitor, significantly reduced the ex vivo production of interferon-gamma by splenic Th1 cells without affecting interleukin-4 production by Th2 cells. FR167653 administration from 4-30 weeks of age prevented NOD mice from developing **diabetes** without affecting the severity of insulinitis. Treatment with FR167653 after insulinitis had developed (i.e. from 10-30 weeks of age) also prevented **diabetes**, further suggesting that treatment with the p38 MAPK pathway inhibitor keeps insulinitis benign in NOD mice, partly by inhibiting Th1 immunity. These findings suggest that p38 MAPK is a key mediator that switches insulinitis from benign to destructive in the development of **type 1 diabetes**.

L22 ANSWER 13 OF 23 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN

ACCESSION NUMBER: 2004:70086 BIOSIS

DOCUMENT NUMBER: PREV200400070419

TITLE: C-peptide enhances insulin-mediated cell growth and protection against high glucose-induced apoptosis in SH-SY5Y cells.

AUTHOR(S): Li, Zhen-guo; Zhang, Weixian; Sima, Anders A. F. [Reprint Author]

CORPORATE SOURCE: Department of Pathology, Wayne State University, 540 East Canfield Ave., Room 9275, H.G. Scott Hall, Detroit, MI, 48201, USA
asima@med.wayne.edu

SOURCE: Diabetes-Metabolism Research and Reviews,
(September-October 2003) Vol. 19, No. 5, pp. 375-385.
print.
ISSN: 1520-7552 (ISSN print).

DOCUMENT TYPE: Article
LANGUAGE: English
ENTRY DATE: Entered STN: 4 Feb 2004
Last Updated on STN: 4 Feb 2004

AB Background: We have previously reported that C-peptide exerts preventive and therapeutic effects on diabetic neuropathy in type 1 diabetic BB/Wor-rats and that it prevents duration-dependent hippocampal apoptosis in the same animal model. In the present study, we examined human neuroblastoma SH-SY5Y cells to examine whether C-peptide stimulates cell proliferation/neurite outgrowth and whether it has antiapoptotic effects. Methods: For neurite outgrowth, serum-starved cultures were treated with C-peptide and/or insulin or IGF-1. Neurites were visualized with NF-L antibody and measured morphometrically. Cell numbers were determined using an electronic cell counter. Scrambled C-peptide was used as a negative control. For assessment of apoptosis, SH-SY5Y cells were incubated with 100 mM glucose for 24 h, and the effects of C-peptide and/or insulin or IGF-1 were examined. Apoptosis was demonstrated by transferase-mediated dUTP nick-end labeling (TUNEL)/4,6-diamidino-2-phenylindole (DAPI) stainings, flow cytometry and changes in the expression of Bcl2. Activation of insulin signaling intermediaries was determined by Western blots. Translocation of NF-kappaB was demonstrated immunocytochemically. Results: C-peptide but not scrambled C-peptide stimulated cell proliferation and neurite outgrowth. In the presence of 4 nM insulin, 3 nM C-peptide significantly increased autophosphorylation of the insulin receptor (IR) but not that of the insulin-like growth factor 1 receptor (IGF-1R). It stimulated phosphoinositide 3-kinase (PI-3 kinase) and p38 mitogen-activated protein (MAP) kinase activation, enhanced the expression and translocation of nuclear factor-kappaB (NF-kappaB), promoted the expression of Bcl2 and reduced c-jun N-terminal kinase (JNK) phosphorylation in excess of that of insulin alone. Conclusions: C-peptide in the presence of insulin exerts synergistic effects on cell proliferation, neurite outgrowth and has in the presence of insulin an antiapoptotic effect on high glucose-induced apoptosis but less so on hyperosmolar-induced apoptosis. These effects are likely to be mediated via interactions with the insulin signaling pathway.

L22 ANSWER 14 OF 23 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on
STN

ACCESSION NUMBER: 2003:471342 BIOSIS
DOCUMENT NUMBER: PREV200300471342
TITLE: "The matrix unloaded": Implications for cytokine signaling
in islets?.
AUTHOR(S): Lowe, William L. Jr. [Reprint Author]
CORPORATE SOURCE: Division of Endocrinology, Metabolism, and Molecular
Medicine, Northwestern University, Feinberg School of
Medicine, 303 East Chicago Avenue, Tarry 15-703, Chicago,
IL, 60611, USA
wlowe@northwestern.edu
SOURCE: Endocrinology, (October 2003) Vol. 144, No. 10, pp.
4262-4263. print.
CODEN: ENDOAO. ISSN: 0013-7227.
DOCUMENT TYPE: Article
General Review; (Literature Review)
LANGUAGE: English
ENTRY DATE: Entered STN: 15 Oct 2003
Last Updated on STN: 15 Oct 2003

L22 ANSWER 15 OF 23 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on
STN

ACCESSION NUMBER: 2003:459325 BIOSIS
DOCUMENT NUMBER: PREV200300459325
TITLE: p38 MAPK inhibitor prevents
the development of **type-1**
diabetes and alleviates hyperglycemia in NOD mice:
A preliminary report.
AUTHOR(S): Medicherla, Satyanarayana [Reprint Author]; Protter, Andrew
[Reprint Author]; Mangadu, Ruban [Reprint Author]; Almirez,
Ramona [Reprint Author]; Ma, Jing [Reprint Author]; Dugar,
Sundeeep [Reprint Author]; Mavunkel, Babu [Reprint Author];
Perumattam, John [Reprint Author]; Schreiner, George
[Reprint Author]
CORPORATE SOURCE: Sunnyvale, CA, USA
SOURCE: Diabetes, (2003) Vol. 52, No. Supplement 1, pp. A126-A127.
print.
Meeting Info.: 63rd Scientific Sessions of the American
Diabetes Association. New Orleans, LA, USA. June 13-17,
2003. American Diabetes Association.
ISSN: 0012-1797 (ISSN print).
DOCUMENT TYPE: Conference; (Meeting)
Conference; Abstract; (Meeting Abstract)
LANGUAGE: English
ENTRY DATE: Entered STN: 8 Oct 2003
Last Updated on STN: 8 Oct 2003

L22 ANSWER 16 OF 23 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on
STN

ACCESSION NUMBER: 2003:104749 BIOSIS
DOCUMENT NUMBER: PREV200300104749
TITLE: Methylglyoxal-bovine serum albumin stimulates tumor
necrosis factor alpha secretion in RAW 264.7 cells through
activation of mitogen-activating protein kinase, nuclear
factor kappaB and intracellular reactive oxygen species
formation.
AUTHOR(S): Fan, X.; Subramaniam, R.; Weiss, M. F.; Monnier, V. M.
[Reprint Author]
CORPORATE SOURCE: Department of Biochemistry, Institute of Pathology, Case
Western Reserve University, 2085 Adelbert Road, Cleveland,
OH, 41106, USA
vmm3@po.cwru.edu
SOURCE: Archives of Biochemistry and Biophysics, (January 15 2003)
Vol. 409, No. 2, pp. 274-286. print.
ISSN: 0003-9861 (ISSN print).
DOCUMENT TYPE: Article
LANGUAGE: English
ENTRY DATE: Entered STN: 19 Feb 2003
Last Updated on STN: 19 Feb 2003

AB Accumulating evidence suggests that the pathophysiology of
diabetes is analogous to chronic inflammatory states. Circulating
levels of inflammatory cytokines such as IL-6 and tumor necrosis factor
alpha (TNFalpha) are increased in both type 1 and type 2 **diabetes**
. TNFalpha plays an important role in the pathogenesis of insulin
resistance in type 2 **diabetes**. However, the reason for this
increase remains unclear. Levels of the dicarbonyl methylglyoxal (MGO)
are elevated in diabetic plasma and MGO-modified bovine serum albumin
(MGO-BSA) can trigger cellular uptake of TNF. Therefore we tested the
hypothesis that MGO-modified proteins may cause TNFalpha secretion in

macrophage-like RAW 264.7 cells. Treatment of cells with MGO-BSA induced TNF α release in a dose-dependent manner. MGO-modified ribonuclease A and chicken egg ovalbumin had similar effects. Cotreatment of cells with antioxidant reagent N-acetylcysteine (NAC) inhibited MGO-BSA-induced TNF α secretion. MGO-BSA stimulated the simultaneous activation of p44/42 and p38 mitogen-activated protein kinase. PD98059, a selective MEK inhibitor, inhibited MGO-BSA-induced TNF α release as well as ERK phosphorylation. Pretreatment of cells with NAC also resulted in inhibition of MGO-BSA-induced ERK phosphorylation. MGO-BSA induced dose-dependent NF κ B activation as shown by electrophoresis mobility shift assay. The MGO-BSA-induced NF κ B activation was prevented in the presence of PD98059, NAC, and parthenolide, a selective inhibitor of NF κ B. Furthermore, the NF κ B inhibitor parthenolide suppressed MGO-BSA-induced TNF α secretion. Confocal microscopy using dichlorofluorescein to demonstrate intracellular reactive oxygen species (ROS) showed that MGO-BSA produced more ROS compared with native BSA. MGO-BSA could also stimulate protein kinase C (PKC) translocation to the cell membrane, considered a key signaling pathway in **diabetes**. However, there was no evidence that PKC was involved in TNF α release based on inhibition by calphostin C and staurosporine. Our findings suggest that the presence of chronically elevated levels of MGO-modified bovine serum albumin may contribute to elevated levels of TNF α in **diabetes**.

L22 ANSWER 17 OF 23 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN

ACCESSION NUMBER: 2003:86411 BIOSIS

DOCUMENT NUMBER: PREV200300086411

TITLE: Are oxidative stress-activated signaling pathways mediators of insulin resistance and beta-cell dysfunction?

AUTHOR(S): Evans, Joseph L. [Reprint Author]; Goldfine, Ira D.; Maddux, Betty A.; Grodsky, Gerold M.

CORPORATE SOURCE: Medical Research Institute, 444 De Haro St., Suite 209, San Francisco, CA, 94107-2347, USA
jevansphd@earthlink.net

SOURCE: Diabetes, (January 2003) Vol. 52, No. 1, pp. 1-8. print.
ISSN: 0012-1797 (ISSN print).

DOCUMENT TYPE: Article
General Review; (Literature Review)

LANGUAGE: English

ENTRY DATE: Entered STN: 6 Feb 2003

Last Updated on STN: 6 Feb 2003

AB In both type 1 and type 2 **diabetes**, diabetic complications in target organs arise from chronic elevations of glucose. The pathogenic effect of high glucose, possibly in concert with fatty acids, is mediated to a significant extent via increased production of reactive oxygen species (ROS) and reactive nitrogen species (RNS) and subsequent oxidative stress. ROS and RNS directly oxidize and damage DNA, proteins, and lipids. In addition to their ability to directly inflict damage on macromolecules, ROS and RNS indirectly induce damage to tissues by activating a number of cellular stress-sensitive pathways. These pathways include nuclear factor- κ B, p38 mitogen-activated protein kinase, NH₂-terminal Jun kinases/stress-activated protein kinases, hexosamines, and others. In addition, there is evidence that in type 2 **diabetes**, the activation of these same pathways by elevations in glucose and free fatty acid (FFA) levels leads to both insulin resistance and impaired insulin secretion. Therefore, we propose here that the hyperglycemia-induced, and possibly FFA-induced, activation of stress pathways plays a key role in the development of not only the late complications in type 1 and type 2 **diabetes**, but also the

insulin resistance and impaired insulin secretion seen in type 2 diabetes.

L22 ANSWER 18 OF 23 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN

ACCESSION NUMBER: 2002:574874 BIOSIS

DOCUMENT NUMBER: PREV200200574874

TITLE: p38 and activating transcription factor-2 involvement in osteoblast osmotic response to elevated extracellular glucose.

AUTHOR(S): Zayzafoon, Majd; Botolin, Sergiu; McCabe, Laura R. [Reprint author]

CORPORATE SOURCE: Dept. of Physiology, Michigan State University, 2201 Biomedical and Physical Sciences Bldg., East Lansing, MI, 48824, USA
mccabel@msu.edu

SOURCE: Journal of Biological Chemistry, (October 4, 2002) Vol. 277, No. 40, pp. 37212-37218. print.
CODEN: JBCHA3. ISSN: 0021-9258.

DOCUMENT TYPE: Article

LANGUAGE: English

ENTRY DATE: Entered STN: 7 Nov 2002

Last Updated on STN: 7 Nov 2002

AB Poorly controlled or untreated **type I diabetes** mellitus is characterized by hyperglycemia and is associated with decreased bone mass and osteoporosis. We have demonstrated that osteoblasts are sensitive to hyperglycemia-associated osmotic stress and respond to elevated extracellular glucose or mannitol by increasing c-jun and collagen I expression. To determine whether MAPKs are involved in this response, MC3T3-E1 osteoblasts were treated with 16.5 mM glucose, mannitol, or contrast dye for 1 h. Immunoblotting of phosphorylated p38 demonstrated activation of p38 MAPK by hyperosmotic stress in vitro and in vivo. Activation peaked at 20 min, remained detectable after 24 h, and was protein kinase C-independent. Activating transcription factor-2 (ATF-2) activation followed the same pattern as phospho-p38. Transactivation of cAMP response element (CRE)- and c-jun promoter (containing a CRE-like element)-reporter constructs increased following hyperosmotic treatment. SB 203580 (a **p38 MAPK inhibitor**) blocked ATF-2 phosphorylation, CRE transactivation, and c-jun promoter activation. Hyperosmotic activation of collagen I promoter activity was also inhibited by SB 203580, consistent with the involvement of c-jun in collagen I up-regulation. Therefore, we propose that hyperglycemia-induced increases in p38 MAPK activity and ATF-2 phosphorylation contribute to CRE activation and modulation of c-jun and collagen I expression in osteoblasts.

L22 ANSWER 19 OF 23 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN

ACCESSION NUMBER: 2002:113519 BIOSIS

DOCUMENT NUMBER: PREV200200113519

TITLE: Heme oxygenase-1 protects pancreatic beta cells from apoptosis caused by various stimuli.

AUTHOR(S): Tobiasch, Edda; Gunther, Lukas; Bach, Fritz H. [Reprint author]

CORPORATE SOURCE: Immunobiology Research Center, Beth Israel Deaconess Medical Center, Harvard Medical School, 99 Brookline Ave, Boston, MA, 02115, USA
fritzhbach@aol.com

SOURCE: Journal of Investigative Medicine, (November, 2001) Vol. 49, No. 6, pp. 566-571. print.

ISSN: 1081-5589.

DOCUMENT TYPE: Article
LANGUAGE: English
ENTRY DATE: Entered STN: 30 Jan 2002
Last Updated on STN: 26 Feb 2002

AB Background: Several problems can occur after allogeneic islet transplantation: primary nonfunction, rejection, and the recurrence of autoimmune disease, which involve attack by the recipient's cytokines, T cells, natural killer cells, and monocytes on the donor's beta cells, which leads to beta-cell destruction. Recent studies have revealed that loss of transplanted islets is caused mainly by apoptosis. Heme oxygenase-1 (HO-1) is one of the antiapoptotic genes up-regulated under stress conditions. The aim of this work was to investigate any mechanisms of HO-1-mediated protection of beta cells from apoptosis. Methods: Apoptosis was assessed by comparison of viable transfected cells with and without apoptotic stimuli, and with and without HO-1 overexpression. Activation and function of p38 mitogen-activated protein kinase were determined using the specific inhibitor SB203580. Results: We have shown that HO-1 mediates antiapoptotic effects in beta cells. The percentage of apoptotic cells after stimulation with tumor necrosis factor alpha decreased from 75% without HO-1 to 5% when HO-1 was overexpressed. Our data indicate that HO-1 acts as a signal terminator of tumor necrosis factor alpha-induced apoptosis by modulation of the p38 mitogen-activated protein kinase pathway. Conclusions: Profound cell stress that occurs in islets after transplantation, as well as at the onset of **diabetes**, results in beta-cell loss through apoptosis. Protection of beta cells by HO-1 improves their survival in vitro after various proapoptotic stimuli, suggesting that HO-1 suppresses one or several signaling pathways leading to apoptosis. We hypothesize that our in vitro findings can be extrapolated to the in vivo situation, and we propose that expression of HO-1 in islets may illuminate a valuable new approach to improving **diabetes** treatment.

L22 ANSWER 20 OF 23 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN

ACCESSION NUMBER: 2001:442256 BIOSIS
DOCUMENT NUMBER: PREV200100442256
TITLE: Activation of renal cortical PI 3-kinase, Akt (PKB) and Erk-1/-2 type MAP kinase but not of p38 MAP kinase, is associated with renal hypertrophy in mice with **type 1 diabetes**.
AUTHOR(S): Faulkner, Jennifer L. [Reprint author]; Gadre, Swarupa [Reprint author]; Senthil, Duraisamy [Reprint author]; Abboud, Hanna E. [Reprint author]; Choudhury, Goutam Ghosh [Reprint author]; Kasinath, Balakuntalam S. [Reprint author]
CORPORATE SOURCE: San Antonio, TX, USA
SOURCE: Diabetes, (June, 2001) Vol. 50, No. Supplement 2, pp. A174. print.
Meeting Info.: 61st Scientific Sessions of the American Diabetes Association. Philadelphia, Pennsylvania, USA. June 22-26, 2001. American Diabetes Association.
CODEN: DIAEAS. ISSN: 0012-1797.
DOCUMENT TYPE: Conference; (Meeting)
Conference; Abstract; (Meeting Abstract)
Conference; (Meeting Poster)
LANGUAGE: English
ENTRY DATE: Entered STN: 19 Sep 2001
Last Updated on STN: 22 Feb 2002

L22 ANSWER 21 OF 23 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN

ACCESSION NUMBER: 2001:245222 BIOSIS
DOCUMENT NUMBER: PREV200100245222
TITLE: Hyperglycemia-dependent, agonist-selective impairment of p38 MAPK activation by diabetic PMN.
AUTHOR(S): McManus, L. M. [Reprint author]; Bloodworth, R. C. [Reprint author]; Ghosh, P. M. [Reprint author]; Pinckard, R. N. [Reprint author]
CORPORATE SOURCE: UTHSCSA, San Antonio, TX, 78229, USA
SOURCE: FASEB Journal, (March 8, 2001) Vol. 15, No. 5, pp. A944. print.
Meeting Info.: Annual Meeting of the Federation of American Societies for Experimental Biology on Experimental Biology 2001. Orlando, Florida, USA. March 31-April 04, 2001.
CODEN: FAJOEC. ISSN: 0892-6638.
DOCUMENT TYPE: Conference; (Meeting)
Conference; Abstract; (Meeting Abstract)
LANGUAGE: English
ENTRY DATE: Entered STN: 23 May 2001
Last Updated on STN: 19 Feb 2002

AB Hyperglycemia-associated infections in **diabetes** reflect a dysfunction in neutrophilic polymorphonuclear leukocytes (PMN). Previously, we observed that PMN from poorly controlled diabetics have reduced functional responsiveness to those agonists which utilize G-protein-coupled receptors (GPCR). The basis for this agonist-selective impairment is unknown but likely reflects altered signal transduction. The present study examined the activation of p38 MAPK in PMN from diabetic (type 1) or normal subjects in relation to glycosylated hemoglobin (HgA1c). Isolated PMN were stimulated with FMLP (100 nM), PAF (1 nM), C5a (10 nM), LTB4 (10 nM), IL-8 (10 nM) or PMA (65 nM); the levels of total or phosphorylated p38 MAPK in cell lysates were determined by Western blotting. Despite the fact that total p38 MAPK was comparable in diabetic and normal PMN, p38 MAPK phosphorylation was markedly reduced in diabetic PMN after stimulation via GPCR. Importantly, this reduction occurred in parallel with a decrease in the level of glycemic control, i.e., as the level of HgA1c was increased, the extent of p38 MAPK activation was decreased. In contrast to the reduction of GPCR-induced p38 MAPK activation in diabetic PMN, the activation of p38 MAPK initiated by PMA was similar to that of normal PMN. This suggests that altered signal transduction in diabetic PMN does not include deficits either at or downstream of PKC activation. In summary, the glycemia-dependent impairment of p38 MAPK activation by diabetic PMN affects diverse agonists which utilize GPCR and likely involves events upstream of PKC activation.

L22 ANSWER 22 OF 23 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN

ACCESSION NUMBER: 2000:438872 BIOSIS
DOCUMENT NUMBER: PREV200000438872
TITLE: Expression of the monocyte chemoattractant protein-1 (MCP-1) in rat and human islet cells and in pre-diabetic NOD mice.
AUTHOR(S): Chen, M.-C. [Reprint author]; Proost, P.; Gysemans, C.; Mathieu, C.; Eizirik, D. L. [Reprint author]
CORPORATE SOURCE: Gene Expression Unit, Diabetes Research Center, Vrije Universiteit Brussel, Brussel, Belgium
SOURCE: Diabetologia, (August, 2000) Vol. 43, No. Supplement 1, pp. A76. print.
Meeting Info.: 36th Annual Meeting of the European Association for the Study of Diabetes. Jerusalem, Israel.

September 17-21, 2000. European Association for the Study of Diabetes.

CODEN: DBTGAI. ISSN: 0012-186X.

DOCUMENT TYPE:

Conference; (Meeting)

Conference; Abstract; (Meeting Abstract)

LANGUAGE:

English

ENTRY DATE:

Entered STN: 18 Oct 2000

Last Updated on STN: 10 Jan 2002

L22 ANSWER 23 OF 23 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights reserved on STN

ACCESSION NUMBER: 2002364103 EMBASE

TITLE: p38 and activating transcription factor-2 involvement in osteoblast osmotic response to elevated extracellular glucose.

AUTHOR: Zayzafoon M.; Botolin S.; McCabe L.R.

CORPORATE SOURCE: L.R. McCabe, Dept. of Physiology, Michigan State University, 2201 Biomed./Physical Sciences Bldg., East Lansing, MI 48824, United States. mccabel@msu.edu

SOURCE: Journal of Biological Chemistry, (4 Oct 2002) Vol. 277, No. 40, pp. 37212-37218. .

Refs: 73

ISSN: 0021-9258 CODEN: JBCHA3

COUNTRY:

United States

DOCUMENT TYPE:

Journal; Article

FILE SEGMENT:

030 Pharmacology

037 Drug Literature Index

LANGUAGE:

English

SUMMARY LANGUAGE:

English

ENTRY DATE:

Entered STN: 31 Oct 2002

Last Updated on STN: 31 Oct 2002

AB Poorly controlled or untreated **type I diabetes**

mellitus is characterized by hyperglycemia and is associated with decreased bone mass and osteoporosis. We have demonstrated that osteoblasts are sensitive to hyperglycemia-associated osmotic stress and respond to elevated extracellular glucose or mannitol by increasing c-jun and collagen I expression. To determine whether MAPKs are involved in this response, MC3T3-E1 osteoblasts were treated with 16.5 mM glucose, mannitol, or contrast dye for 1 h. Immunoblotting of phosphorylated p38 demonstrated activation of p38 MAPK by hyperosmotic stress in vitro and in vivo. Activation peaked at 20 min, remained detectable after 24 h, and was protein kinase C-independent. Activating transcription factor-2 (ATF-2) activation followed the same pattern as phospho-p38. Transactivation of cAMP response element (CRE)- and c-jun promoter (containing a CRE-like element)-reporter constructs increased following hyperosmotic treatment. SB 203580 (a **p38 MAPK inhibitor**) blocked ATF-2 phosphorylation, CRE transactivation, and c-jun promoter activation. Hyperosmotic activation of collagen I promoter activity was also inhibited by SB 203580, consistent with the involvement of c-jun in collagen I up-regulation. Therefore, we propose that hyperglycemia-induced increases in p38 MAPK activity and ATF-2 phosphorylation contribute to CRE activation and modulation of c-jun and collagen I expression in osteoblasts.